



US011338044B2

(12) **United States Patent**
Guild et al.

(10) **Patent No.:** **US 11,338,044 B2**
(45) **Date of Patent:** **May 24, 2022**

(54) **LIPID NANOPARTICLE COMPOSITIONS AND METHODS FOR MRNA DELIVERY**

(71) Applicant: **Translate Bio, Inc.**, Lexington, MA (US)

(72) Inventors: **Braydon Charles Guild**, Lexington, MA (US); **Frank DeRosa**, Lexington, MA (US); **Michael Heartlein**, Lexington, MA (US)

(73) Assignee: **Translate Bio, Inc.**, Lexington, MA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/556,974**

(22) Filed: **Dec. 20, 2021**

(65) **Prior Publication Data**

US 2022/0105201 A1 Apr. 7, 2022

Related U.S. Application Data

(62) Division of application No. 17/509,674, filed on Oct. 25, 2021, now Pat. No. 11,291,734, which is a division of application No. 17/341,016, filed on Jun. 7, 2021, now Pat. No. 11,185,595, which is a division of application No. 17/099,462, filed on Nov. 16, 2020, now Pat. No. 11,052,159, which is a division of application No. 16/931,231, filed on Jul. 16, 2020, now Pat. No. 10,888,626, which is a division of application No. 16/681,565, filed on Nov. 12, 2019, now abandoned, which is a division of application No. 16/502,672, filed on Jul. 3, 2019, now Pat. No. 10,507,249, which is a division of application No. 16/286,400, filed on Feb. 26, 2019, now Pat. No. 10,350,303, which is a division of application No. 16/233,031, filed on Dec. 26, 2018, now Pat. No. 10,413,618, which is a division of application No. 15/482,117, filed on Apr. 7, 2017, now Pat. No. 10,238,754, which is a division of application No. 14/124,608, filed as application No. PCT/US2012/041724 on Jun. 8, 2012, now abandoned.

(60) Provisional application No. 61/494,881, filed on Jun. 8, 2011.

(51) **Int. Cl.**
B82Y 5/00 (2011.01)
A61K 38/18 (2006.01)
A61K 48/00 (2006.01)
A61K 9/19 (2006.01)
A61K 9/00 (2006.01)
A61K 9/127 (2006.01)
A61K 9/51 (2006.01)
A61K 38/47 (2006.01)
A61K 38/48 (2006.01)
C12N 15/85 (2006.01)
A61K 31/713 (2006.01)

A61K 38/57 (2006.01)
A61K 31/7105 (2006.01)
A61K 31/7115 (2006.01)
C07K 14/505 (2006.01)
C07K 14/81 (2006.01)
C12N 9/40 (2006.01)
C12N 9/64 (2006.01)
C12N 15/52 (2006.01)

(52) **U.S. Cl.**
CPC **A61K 48/0008** (2013.01); **A61K 9/0019** (2013.01); **A61K 9/0073** (2013.01); **A61K 9/1271** (2013.01); **A61K 9/1272** (2013.01); **A61K 9/5123** (2013.01); **A61K 31/713** (2013.01); **A61K 38/1816** (2013.01); **A61K 38/47** (2013.01); **A61K 38/4846** (2013.01); **A61K 38/57** (2013.01); **A61K 48/0075** (2013.01); **A61K 48/0091** (2013.01); **C07K 14/505** (2013.01); **C07K 14/8125** (2013.01); **C12N 9/2465** (2013.01); **C12N 9/644** (2013.01); **C12N 15/52** (2013.01); **C12N 15/85** (2013.01); **C12Y 302/01022** (2013.01); **C12Y 304/21022** (2013.01); **A61K 48/00** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,647,121 A 7/1953 Jacoby
2,717,909 A 9/1955 Kosmin
2,819,718 A 1/1958 Goldman
2,844,629 A 7/1958 William et al.
3,614,955 A 10/1971 Mirowski
3,656,185 A 4/1972 Carpentier
3,805,301 A 4/1974 Liebig
3,945,052 A 3/1976 Liebig
3,995,623 A 12/1976 Blake et al.
4,013,507 A 3/1977 Rembaum

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2518132 A1 3/2006
CA 2807552 A1 2/2012

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 60/083,294.

(Continued)

Primary Examiner — Robert M Kelly

(74) *Attorney, Agent, or Firm* — Proskauer Rose LLP

(57) **ABSTRACT**

Disclosed herein are compositions and methods for modulating the production of a protein in a target cell. The compositions and methods disclosed herein are capable of ameliorating diseases associated with protein or enzyme deficiencies.

21 Claims, 28 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

4,072,146 A	2/1978	Howes	5,595,756 A	1/1997	Bally et al.
4,096,860 A	6/1978	McLaughlin	5,607,385 A	3/1997	Francischelli et al.
4,099,528 A	7/1978	Sorenson et al.	5,609,624 A	3/1997	Kalis
4,106,129 A	8/1978	Carpentier et al.	5,610,283 A	3/1997	Buechler
4,134,402 A	1/1979	Mahurkar	5,614,548 A	3/1997	Piantadosi et al.
4,140,126 A	2/1979	Choudhury	5,626,869 A	5/1997	Nyqvist et al.
4,180,068 A	12/1979	Jacobsen et al.	5,631,018 A	5/1997	Zalipsky et al.
4,182,833 A	1/1980	Hicks	5,677,124 A	10/1997	DuBois et al.
4,227,533 A	10/1980	Godfrey	5,693,088 A	12/1997	Lazarus
4,284,459 A	8/1981	Patel et al.	5,697,953 A	12/1997	Kroll et al.
4,308,085 A	12/1981	Horhold et al.	5,700,437 A	12/1997	Fujii et al.
4,323,525 A	4/1982	Bornat	5,705,188 A	1/1998	Junichi et al.
4,335,723 A	6/1982	Patel	5,705,385 A	1/1998	Bally et al.
4,339,369 A	7/1982	Hicks et al.	5,736,573 A	4/1998	Galat
4,355,426 A	10/1982	MacGregor	5,744,335 A	4/1998	Wolff et al.
4,375,817 A	3/1983	Engle et al.	5,772,694 A	6/1998	Bokros et al.
4,385,631 A	5/1983	Uthmann	5,776,165 A	7/1998	Ripart
4,401,472 A	8/1983	Gerber	5,776,747 A	7/1998	Schinstine et al.
4,406,656 A	9/1983	Hattler et al.	5,783,383 A	7/1998	Kondo et al.
4,475,972 A	10/1984	Wong	5,783,566 A	7/1998	Mislick
4,530,113 A	7/1985	Matterson	5,844,107 A	12/1998	Hanson et al.
4,550,447 A	11/1985	Seiler, Jr. et al.	5,874,105 A	2/1999	Watkins et al.
4,562,596 A	1/1986	Kornberg	5,885,613 A	3/1999	Holland et al.
4,568,329 A	2/1986	Mahurkar	5,910,168 A	6/1999	Myers et al.
4,571,241 A	2/1986	Christopher	5,916,208 A	6/1999	Luther et al.
4,601,718 A	7/1986	Possis et al.	5,965,434 A	10/1999	Wolff et al.
4,647,416 A	3/1987	Seiler, Jr. et al.	5,976,567 A	11/1999	Wheeler et al.
4,662,382 A	5/1987	Shuetz et al.	5,976,569 A	11/1999	Milstein
4,701,162 A	10/1987	Rosenberg	5,981,501 A	11/1999	Wheeler et al.
4,710,169 A	12/1987	Christopher	6,055,454 A	4/2000	Heemels
4,720,517 A	1/1988	Ravichandran et al.	6,067,471 A	5/2000	Warren
4,737,323 A	4/1988	Martin et al.	6,077,835 A	6/2000	Hanson et al.
4,762,915 A	8/1988	Kung et al.	6,090,384 A	7/2000	Ra et al.
4,782,836 A	11/1988	Alt	6,093,392 A	7/2000	High et al.
4,856,521 A	8/1989	Irnich	6,096,070 A	8/2000	Ragheb et al.
4,860,751 A	8/1989	Callaghan	6,096,075 A	8/2000	Bokros et al.
4,878,908 A	11/1989	Martin et al.	6,120,799 A	9/2000	McDonald et al.
4,892,540 A	1/1990	Vallana	6,147,055 A	11/2000	Hobart et al.
4,897,355 A	1/1990	Eppstein et al.	6,152,955 A	11/2000	KenKnight et al.
4,920,016 A	4/1990	Allen et al.	6,165,763 A	12/2000	Brown et al.
4,946,683 A	8/1990	Forssen	6,169,923 B1	1/2001	Kroll
4,946,857 A	8/1990	Kanehira et al.	6,176,877 B1	1/2001	Buchanan et al.
4,960,409 A	10/1990	Catalano	6,204,297 B1	3/2001	Tracy et al.
4,966,945 A	10/1990	Drawer et al.	6,210,892 B1	4/2001	Bennett et al.
5,024,671 A	6/1991	Tu et al.	6,214,804 B1	4/2001	Feigner et al.
5,025,005 A	6/1991	Nomura et al.	6,271,208 B1	8/2001	Bischoff
5,047,540 A	9/1991	Kamata et al.	6,271,209 B1	8/2001	Smith et al.
5,101,824 A	4/1992	Lekholm	6,287,591 B1	9/2001	Semple et al.
5,104,399 A	4/1992	Lazarus	6,299,604 B1	10/2001	Ragheb et al.
5,116,360 A	5/1992	Pinchuk et al.	6,335,199 B1	1/2002	Bischoff et al.
5,138,067 A	8/1992	Kamata et al.	6,358,278 B1	3/2002	Brendzel et al.
5,151,105 A	9/1992	Kwan-Gett	6,370,434 B1	4/2002	Zhang et al.
5,171,678 A	12/1992	Behr et al.	6,371,983 B1	4/2002	Lane
5,176,661 A	1/1993	Evard et al.	6,417,326 B1	7/2002	Cullis et al.
5,194,654 A	3/1993	Hostetler et al.	6,485,726 B1	11/2002	Blumberg et al.
5,200,395 A	4/1993	Eto et al.	6,534,484 B1	3/2003	Wheeler et al.
5,223,263 A	6/1993	Hostetler et al.	6,586,410 B1	7/2003	Wheeler et al.
5,261,419 A	11/1993	Osypka	6,670,178 B1	12/2003	Selden et al.
5,264,618 A	11/1993	Feigner et al.	6,696,424 B1	2/2004	Wheeler
5,279,833 A	1/1994	Rose	6,733,777 B2	5/2004	Erbacher et al.
5,282,824 A	2/1994	Gianturco	6,734,171 B1	5/2004	Saravolac et al.
5,284,491 A	2/1994	Sutton et al.	6,743,823 B1	6/2004	Summar et al.
5,300,022 A	4/1994	Klapper et al.	6,756,055 B2	6/2004	McDonald et al.
5,314,430 A	5/1994	Bardy	6,790,838 B2	9/2004	Alison et al.
5,330,768 A	7/1994	Park et al.	6,815,432 B2	11/2004	Wheeler et al.
5,334,761 A	8/1994	Gebeyehu et al.	6,821,530 B2	11/2004	Koob et al.
5,395,619 A	3/1995	Zalipsky et al.	6,835,395 B1	12/2004	Semple et al.
5,405,363 A	4/1995	Kroll et al.	6,858,224 B2	2/2005	Wheeler et al.
5,405,379 A	4/1995	Lane	6,858,225 B2	2/2005	Semple et al.
5,455,352 A	10/1995	Huellmann et al.	6,887,665 B2	5/2005	Trulson et al.
5,464,924 A	11/1995	Silvis et al.	6,998,115 B2	2/2006	Langer et al.
5,503,852 A	4/1996	Steiner et al.	7,022,214 B2	4/2006	Olech
5,528,023 A	6/1996	Butturini et al.	7,067,697 B2	6/2006	Gao
5,552,155 A	9/1996	Bailey et al.	7,084,303 B2	8/2006	Watanabe et al.
5,580,859 A	12/1996	Feigner et al.	7,341,738 B2	3/2008	Semple et al.
			7,422,902 B1	9/2008	Wheeler et al.
			7,427,394 B2	9/2008	Anderson et al.
			7,507,859 B2	3/2009	Grinstaff et al.
			7,556,684 B2	7/2009	Bury et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

7,803,397 B2	9/2010	Heyes et al.	9,428,535 B2	8/2016	de Fougerolles et al.
7,888,067 B2	2/2011	Lin et al.	9,428,751 B2	8/2016	MacDonald et al.
7,901,708 B2	3/2011	MacLachlan et al.	9,464,124 B2	10/2016	Bancel et al.
7,972,435 B2	7/2011	Bury et al.	9,492,386 B2	11/2016	MacLachlan et al.
8,021,686 B2	9/2011	Semple et al.	9,504,651 B2	11/2016	MacLachlan et al.
8,071,082 B2	12/2011	Zugates et al.	9,504,734 B2	11/2016	Bancel et al.
8,101,741 B2	1/2012	MacLachlan et al.	9,518,272 B2	12/2016	Yaworksi et al.
8,106,022 B2	1/2012	Manoharan et al.	9,572,874 B2	2/2017	Fotin-Mleczeck et al.
8,158,601 B2	4/2012	Chen et al.	9,572,896 B2	2/2017	Bancel et al.
8,188,263 B2	5/2012	MacLachlan et al.	9,572,897 B2	2/2017	Bancel et al.
RE43,612 E	8/2012	Anderson et al.	9,587,003 B2	3/2017	Bancel et al.
8,236,943 B2	8/2012	Lee et al.	9,616,084 B2	4/2017	Mutzke
8,278,036 B2	10/2012	Kariko et al.	9,623,095 B2	4/2017	Kallen et al.
8,287,849 B2	10/2012	Langer et al.	9,636,301 B2	5/2017	Weber
8,329,070 B2	12/2012	MacLachlan et al.	9,669,089 B2	6/2017	Thess et al.
8,450,298 B2	5/2013	Mahon et al.	9,675,668 B2	6/2017	Bancel et al.
8,450,467 B2	5/2013	Manoharan et al.	9,682,139 B2	6/2017	Manoharan et al.
8,513,403 B2	8/2013	MacLachlan et al.	9,683,233 B2	6/2017	Thess
8,557,231 B2	10/2013	Langer et al.	9,688,729 B2	6/2017	Kramps et al.
8,562,966 B2	10/2013	Zugates et al.	10,238,754 B2	3/2019	Guild et al.
8,569,256 B2	10/2013	Heyes et al.	10,350,303 B1	7/2019	Guild et al.
8,652,512 B2	2/2014	Schmehl et al.	10,413,618 B2	9/2019	Guild et al.
8,691,966 B2	4/2014	Kariko et al.	10,507,249 B2	12/2019	Guild et al.
8,710,200 B2	4/2014	Schrum et al.	10,888,626 B2	1/2021	Guild et al.
8,748,089 B2	6/2014	Kariko et al.	11,052,159 B2	7/2021	Guild et al.
8,802,644 B2	8/2014	Chen et al.	2002/0022721 A1	2/2002	Trulson et al.
8,808,681 B2	8/2014	Anderson et al.	2002/0094528 A1	7/2002	Salatsky
8,808,982 B2	8/2014	Dahl et al.	2002/0192651 A1	12/2002	Wheeler et al.
8,822,663 B2	9/2014	Schrum et al.	2002/0192721 A1	12/2002	Rizzuto et al.
8,828,956 B2	9/2014	Manoharan et al.	2002/0193622 A1	12/2002	Watanabe et al.
8,835,108 B2	9/2014	Kariko et al.	2003/0082154 A1	5/2003	Leamon
8,846,348 B2	9/2014	Jendrisak et al.	2003/0083272 A1	5/2003	Wiederholt et al.
8,853,377 B2	10/2014	Guild et al.	2003/0104044 A1	6/2003	Semple et al.
8,859,229 B2	10/2014	Rabinovich et al.	2003/0215395 A1	11/2003	Yu et al.
8,883,202 B2	11/2014	Manoharan et al.	2004/0110709 A1	6/2004	Li et al.
8,936,942 B2	1/2015	Heyes et al.	2004/0132683 A1	7/2004	Feigner et al.
8,969,353 B2	3/2015	Mahon et al.	2004/0224912 A1	11/2004	Dobie et al.
8,999,351 B2	4/2015	Manoharan et al.	2004/0235982 A1	11/2004	Rabasco et al.
8,999,950 B2	4/2015	MacLachlan et al.	2005/0004058 A1	1/2005	Benoit et al.
9,005,930 B2	4/2015	Jendrisak et al.	2005/0008689 A1	1/2005	Semple et al.
9,012,219 B2	4/2015	Kariko et al.	2005/0032730 A1	2/2005	Von Der Mulbe et al.
9,012,498 B2	4/2015	Manoharan et al.	2005/0054026 A1	3/2005	Atsushi et al.
9,018,187 B2	4/2015	Heyes et al.	2005/0059005 A1	3/2005	Tuschl et al.
9,040,256 B2	5/2015	Grunenwald et al.	2005/0059624 A1	3/2005	Hoerr et al.
9,051,567 B2	6/2015	Fitzgerald et al.	2005/0065107 A1	3/2005	Hobart et al.
9,061,021 B2	6/2015	Guild et al.	2005/0069590 A1	3/2005	Buehler et al.
9,061,059 B2	6/2015	Chakraborty et al.	2005/0079212 A1	4/2005	Wheeler et al.
9,074,208 B2	7/2015	MacLachlan et al.	2005/0143332 A1	6/2005	Monahan et al.
9,080,211 B2	7/2015	Grunenwald et al.	2005/0148786 A1	7/2005	Ikeda et al.
9,085,801 B2	7/2015	Grunenwald et al.	2005/0158302 A1	7/2005	Faustman et al.
9,089,604 B2	7/2015	Chakraborty et al.	2005/0244961 A1	11/2005	Short et al.
9,095,552 B2	8/2015	Chakraborty et al.	2005/0250723 A1	11/2005	Hoerr et al.
9,107,886 B2	8/2015	Chakraborty et al.	2006/0008910 A1	1/2006	MacLachlan et al.
9,114,113 B2	8/2015	Chakraborty et al.	2006/0059576 A1	3/2006	Pasinetti et al.
9,181,319 B2	11/2015	Schrum et al.	2006/0069225 A1	3/2006	Wintermantel et al.
9,181,321 B2	11/2015	Heartlein et al.	2006/0083780 A1	4/2006	Heyes et al.
9,186,325 B2	11/2015	Manoharan et al.	2006/0172003 A1	8/2006	Meers et al.
9,186,372 B2	11/2015	de Fougerolles et al.	2006/0204566 A1	9/2006	Smyth-Templeton et al.
9,187,748 B2	11/2015	Geisbert et al.	2006/0216343 A1	9/2006	Panzner et al.
9,192,651 B2	11/2015	Chakraborty et al.	2006/0223939 A1	10/2006	Lange et al.
9,220,682 B2	12/2015	Manoharan et al.	2006/0228404 A1	10/2006	Anderson et al.
9,220,755 B2	12/2015	Chakraborty et al.	2006/0241071 A1	10/2006	Grinstaff et al.
9,220,792 B2	12/2015	Chakraborty et al.	2007/0142628 A1	6/2007	Ghoshal et al.
9,233,141 B2	1/2016	Chakraborty et al.	2007/0172950 A1	7/2007	Wheeler et al.
9,254,311 B2	2/2016	Bancel et al.	2007/0252295 A1	11/2007	Panzner et al.
9,295,689 B2	3/2016	de Fougerolles et al.	2007/0275923 A1	11/2007	Chen et al.
9,301,993 B2	4/2016	Chakraborty et al.	2007/0281336 A1	12/2007	Jendrisak et al.
9,303,079 B2	4/2016	Chakraborty et al.	2008/0145338 A1	6/2008	Anderson et al.
9,334,328 B2	5/2016	Schrum et al.	2008/0160048 A1	7/2008	Fuller
9,345,780 B2	5/2016	Manoharan et al.	2008/0242626 A1	10/2008	Zugates et al.
9,352,042 B2	5/2016	Heyes et al.	2008/0260706 A1	10/2008	Rabinovich et al.
9,352,048 B2	5/2016	Manoharan et al.	2009/0023673 A1	1/2009	Manoharan et al.
9,364,435 B2	6/2016	Yaworski et al.	2009/0093433 A1	4/2009	Wolf et al.
9,394,234 B2	7/2016	Chen et al.	2009/0163705 A1	6/2009	Manoharan et al.
9,404,127 B2	8/2016	Yaworski et al.	2009/0186805 A1	7/2009	Tabor et al.
			2009/0221684 A1	9/2009	Grinstaff et al.
			2009/0263407 A1	10/2009	Dande et al.
			2009/0270481 A1	10/2009	MacLachlan et al.
			2009/0286852 A1	11/2009	Kariko et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2009/0326051	A1	12/2009	Corey et al.	2014/0255468	A1	9/2014	Bancel et al.
2010/0028943	A1	2/2010	Thomas et al.	2014/0275227	A1	9/2014	Hoge et al.
2010/0035249	A1	2/2010	Hayashizaki et al.	2014/0275229	A1	9/2014	Bancel et al.
2010/0036084	A1	2/2010	Langer et al.	2014/0288160	A1	9/2014	Guild et al.
2010/0041152	A1	2/2010	Wheeler et al.	2014/0294937	A1	10/2014	MacLachlan et al.
2010/0047261	A1	2/2010	Hoerr et al.	2014/0294938	A1	10/2014	Guild et al.
2010/0120129	A1	5/2010	Amshey et al.	2014/0294940	A1	10/2014	Guild et al.
2010/0178699	A1	7/2010	Gao et al.	2014/0329884	A1	11/2014	Dong et al.
2010/0189729	A1	7/2010	Hoerr et al.	2014/0343129	A1	11/2014	de Fougerolles et al.
2010/0267806	A1	10/2010	Bumcrot et al.	2014/0363876	A1	12/2014	Jendrisak et al.
2010/0323356	A1	12/2010	Inoue et al.	2015/0004217	A1	1/2015	Guild et al.
2010/0331234	A1	12/2010	Mahon et al.	2015/0005372	A1	1/2015	Hoge et al.
2011/0009641	A1	1/2011	Anderson et al.	2015/0011615	A1	1/2015	Manoharan et al.
2011/0038941	A1	2/2011	Lee et al.	2015/0011633	A1	1/2015	Shorr et al.
2011/0092739	A1	4/2011	Chen et al.	2015/0017211	A1	1/2015	de Fougerolles et al.
2011/0143397	A1	6/2011	Kariko et al.	2015/0038556	A1	2/2015	Heartlein et al.
2011/0200582	A1	8/2011	Baryza et al.	2015/0038558	A1	2/2015	Kariko et al.
2011/0244026	A1	10/2011	Guild et al.	2015/0044277	A1	2/2015	Bancel et al.
2011/0256175	A1	10/2011	Hope et al.	2015/0050354	A1	2/2015	Bouchon et al.
2011/0287435	A1	11/2011	Grunenwald et al.	2015/0051268	A1	2/2015	Bancel et al.
2011/0293703	A1	12/2011	Mahon et al.	2015/0056253	A1	2/2015	Bancel et al.
2011/0311583	A1	12/2011	Manoharan et al.	2015/0064235	A1	3/2015	Bancel et al.
2012/0007803	A1	1/2012	Takatsuka	2015/0064236	A1	3/2015	Bancel et al.
2012/0009222	A1	1/2012	Nguyen et al.	2015/0064242	A1	3/2015	Heyes et al.
2012/0065252	A1	3/2012	Schrum et al.	2015/0064725	A1	3/2015	Schrum et al.
2012/0065358	A1	3/2012	Langer et al.	2015/0086614	A1	3/2015	Bancel et al.
2012/0114831	A1	5/2012	Semple et al.	2015/0110857	A1	4/2015	DeRosa et al.
2012/0128760	A1	5/2012	Manoharan et al.	2015/0110858	A1	4/2015	DeRosa et al.
2012/0129910	A1	5/2012	Thompson et al.	2015/0110859	A1	4/2015	Heartlein et al.
2012/0142756	A1	6/2012	Guild et al.	2015/0119444	A1	4/2015	Manoharan et al.
2012/0195936	A1	8/2012	Rudolph et al.	2015/0119445	A1	4/2015	Manoharan et al.
2012/0237975	A1	9/2012	Schrum et al.	2015/0157565	A1	6/2015	Heartlein et al.
2012/0251560	A1	10/2012	Dahlman et al.	2015/0166465	A1	6/2015	Chen et al.
2012/0251618	A1	10/2012	Schrum et al.	2015/0190515	A1	7/2015	Manoharan et al.
2013/0017223	A1	1/2013	Hope et al.	2015/0191760	A1	7/2015	Jendrisak et al.
2013/0158021	A1	6/2013	Dong et al.	2015/0265708	A1	9/2015	Manoharan et al.
2013/0195967	A1	8/2013	Guild et al.	2015/0267192	A1	9/2015	Heartlein et al.
2013/0259923	A1	10/2013	Chakraborty et al.	2015/0315541	A1	11/2015	Bancel et al.
2013/0259924	A1	10/2013	Bancel et al.	2015/0315584	A1	11/2015	MacDonald et al.
2013/0266640	A1	10/2013	de Fougerolles et al.	2015/0366997	A1	12/2015	Guild et al.
2013/0302401	A1	11/2013	Ma et al.	2016/0095924	A1	4/2016	Hope et al.
2014/0010861	A1	1/2014	Bancel et al.	2016/0114011	A1	4/2016	Bancel et al.
2014/0044772	A1	2/2014	MacLachlan et al.	2016/0115477	A1	4/2016	MacLachlan et al.
2014/0094399	A1	4/2014	Langer et al.	2016/0115483	A1	4/2016	MacLachlan et al.
2014/0105964	A1	4/2014	Chakraborty et al.	2016/0136236	A1	5/2016	Hoge et al.
2014/0105965	A1	4/2014	Chakraborty et al.	2016/0151284	A1	6/2016	Heyes et al.
2014/0147432	A1	5/2014	Chakraborty et al.	2016/0158385	A1	6/2016	Bancel et al.
2014/0147454	A1	5/2014	Chakraborty et al.	2016/0193299	A1	7/2016	de Fougerolles et al.
2014/0148502	A1	5/2014	Bancel et al.	2016/0194368	A1	7/2016	Hoge et al.
2014/0155472	A1	6/2014	Bancel et al.	2016/0194625	A1	7/2016	Hoge et al.
2014/0155473	A1	6/2014	Bancel et al.	2016/0199485	A1	7/2016	Manoharan et al.
2014/0155474	A1	6/2014	Bancel et al.	2016/0213785	A1	7/2016	Manoharan et al.
2014/0155475	A1	6/2014	Bancel et al.	2016/0237108	A1	8/2016	Fraley et al.
2014/0161830	A1	6/2014	Anderson et al.	2016/0237134	A1	8/2016	Hoge et al.
2014/0162897	A1	6/2014	Grunenwald et al.	2016/0250354	A1	9/2016	Manoharan et al.
2014/0171485	A1	6/2014	Bancel et al.	2016/0251681	A1	9/2016	Yaworski et al.
2014/0179771	A1	6/2014	Bancel et al.	2016/0256567	A1	9/2016	Heyes et al.
2014/0186432	A1	7/2014	Chakraborty et al.	2016/0256568	A1	9/2016	Heyes et al.
2014/0193482	A1	7/2014	Chakraborty et al.	2016/0256573	A1	9/2016	de Fougerolles et al.
2014/0194494	A1	7/2014	Bancel et al.	2016/0264971	A1	9/2016	Geisbert et al.
2014/0199371	A1	7/2014	Chakraborty et al.	2016/0264975	A1	9/2016	Schrum et al.
2014/0200261	A1	7/2014	Hoge et al.	2016/0274089	A1	9/2016	Ciufolini et al.
2014/0200262	A1	7/2014	Chakraborty et al.	2016/0304552	A1	10/2016	Roy et al.
2014/0200263	A1	7/2014	Bancel et al.	2016/0317647	A1	11/2016	Ciaramella et al.
2014/0200264	A1	7/2014	Bancel et al.	2016/0317676	A1	11/2016	Hope et al.
2014/0206752	A1	7/2014	Afeyan et al.	2016/0331828	A1	11/2016	Ciaramella et al.
2014/0206753	A1	7/2014	Guild et al.	2016/0348099	A1	12/2016	Roy et al.
2014/0206755	A1	7/2014	Bancel et al.	2016/0354490	A1	12/2016	Roy et al.
2014/0206852	A1	7/2014	Bancel et al.	2016/0354491	A1	12/2016	Roy et al.
2014/0221248	A1	8/2014	Jendrisak et al.	2016/0354492	A1	12/2016	Roy et al.
2014/0221465	A1	8/2014	Bancel et al.	2016/0354493	A1	12/2016	Roy et al.
2014/0227300	A1	8/2014	Chin et al.	2016/0367687	A1	12/2016	Manoharan et al.
2014/0243399	A1	8/2014	Schrum et al.	2016/0367702	A1	12/2016	Hoge et al.
2014/0249208	A1	9/2014	Bancel et al.	2016/0375134	A1	12/2016	Bancel et al.
2014/0255467	A1	9/2014	Bancel et al.	2016/0375137	A9	12/2016	Manoharan et al.
				2017/0000858	A1	1/2017	Fotin-Mleczeck et al.
				2017/0000870	A1	1/2017	Hoerr et al.
				2017/0000871	A1	1/2017	Probst et al.
				2017/0002060	A1	1/2017	Bolen et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2017/0007702 A1 1/2017 Heyes et al.
 2017/0014496 A1 1/2017 Fotin-Mleczek et al.
 2017/0027658 A1 2/2017 Black et al.
 2017/0028059 A1 2/2017 Baumhof et al.
 2017/0029847 A1 2/2017 Thess
 2017/0042814 A1 2/2017 Yaworski et al.
 2017/0056528 A1 3/2017 De Fougerolles et al.
 2017/0056529 A1 3/2017 Thess et al.
 2017/0065675 A1 3/2017 Bancel et al.
 2017/0065727 A1 3/2017 Fotin-Mleczek et al.
 2017/0114378 A1 4/2017 Wochner et al.
 2017/0128549 A1 5/2017 Fotin-Mileczek et al.
 2017/0136131 A1 5/2017 Atanu et al.
 2017/0157268 A1 6/2017 Ansell et al.
 2017/0166905 A1 6/2017 Eberle et al.
 2017/0173128 A1 6/2017 Hoge et al.
 2017/0175129 A1 6/2017 Roy et al.
 2017/0182081 A1 6/2017 Mutzke
 2017/0182150 A1 6/2017 Kallen et al.
 2017/0190661 A1 6/2017 Payne et al.
 2017/0189552 A1 7/2017 Hasenpusch
 2018/0161451 A1 1/2018 Fotin-Mleczek et al.

FOREIGN PATENT DOCUMENTS

CN 1399561 2/2003
 CN 100569877 C 12/2009
 CN 101863544 A 10/2010
 DE 24 30 998 A1 1/1975
 DE 2520814 A1 11/1976
 DE 3728917 A1 3/1989
 EP 6 73 637 A1 9/1995
 EP 0783297 A1 7/1997
 EP 0853123 A1 7/1998
 EP 0959092 A1 11/1999
 EP 2449106 A1 11/1999
 EP 1026253 A2 8/2000
 EP 1519714 B1 4/2005
 EP 1979364 A2 10/2008
 EP 2045251 A1 4/2009
 EP 2338478 B1 6/2011
 EP 2338520 A1 6/2011
 EP 2532649 A1 12/2012
 EP 2578685 A2 4/2013
 EP 2823809 A1 1/2015
 FR 1 378 382 A 11/1964
 FR 2 235 112 A1 1/1975
 GB 1072118 A 6/1967
 GB 1602085 A 11/1981
 JP H07-053535 2/1955
 JP S48-022365 3/1973
 JP S49-127908 A 12/1974
 JP S51-023537 2/1976
 JP 51-125144 11/1976
 JP S52-010847 1/1977
 JP 63-125144 5/1988
 JP 63-154788 6/1988
 JP H09-505593 6/1997
 JP H10-197978 7/1998
 JP 11-005786 A 1/1999
 JP 11-080142 3/1999
 JP 2001-523215 A 11/2001
 JP 2002-167368 A 6/2002
 JP 2003-519199 A 6/2003
 JP 4-108173 B2 6/2008
 JP 2008-247749 10/2008
 JP 50-24216 B2 9/2012
 RU 2248213 C2 3/2005
 WO WO-1990/11092 A1 10/1990
 WO WO-1993/18229 A1 9/1993
 WO WO-1993/18754 A1 9/1993
 WO WO-1995/11004 A1 4/1995
 WO WO-1995/14651 A1 6/1995
 WO WO-1995/27478 A1 10/1995
 WO WO-1996/18372 A2 6/1996

WO WO-1996/26179 A1 8/1996
 WO WO-1996/37211 A1 11/1996
 WO WO-1996/40964 A2 12/1996
 WO WO-1997/46223 A1 12/1997
 WO WO-1998/10748 A1 3/1998
 WO WO-1998/16202 A2 4/1998
 WO WO-1998/51278 A2 11/1998
 WO WO-1999/14346 A2 3/1999
 WO WO-1999/18933 A2 4/1999
 WO WO-2000/03044 A1 1/2000
 WO WO-2000/62813 A2 10/2000
 WO WO-2000/64484 A2 11/2000
 WO WO-2000/69913 A1 11/2000
 WO WO-2001/05375 A1 1/2001
 WO WO-2001/07599 A1 2/2001
 WO WO-2002/00870 A2 1/2002
 WO WO-2002/22709 A1 3/2002
 WO WO-2002/31025 A2 4/2002
 WO WO-2002/34236 A2 5/2002
 WO WO-2002/42317 A2 5/2002
 WO WO-2003/040288 A2 5/2003
 WO WO-2003/070735 A2 8/2003
 WO WO-2004/002453 A1 1/2004
 WO WO-2004/043588 A2 5/2004
 WO WO-2004/048345 A2 6/2004
 WO WO-2004/106411 A2 12/2004
 WO WO-2005/026372 A1 3/2005
 WO WO-2005/028619 A2 3/2005
 WO WO-2005/037226 A2 4/2005
 WO WO2005/120152 A2 12/2005
 WO WO-2005/121348 A1 12/2005
 WO WO-2006/000448 A2 1/2006
 WO WO-2006/016097 A2 2/2006
 WO WO-2006/082088 A1 8/2006
 WO WO-2006/105043 A2 10/2006
 WO WO-2006/138380 A2 12/2006
 WO WO-2007/024708 A2 3/2007
 WO WO-2007/031091 A2 3/2007
 WO WO-2007/120863 A2 10/2007
 WO WO-2007/126386 A1 11/2007
 WO WO-2007/143659 A2 12/2007
 WO WO-2008/011561 A2 1/2008
 WO WO-2008/042973 A2 4/2008
 WO WO-2008/045548 A2 4/2008
 WO WO2008/052770 A2 5/2008
 WO WO-2008/083949 A2 7/2008
 WO WO-2008/113364 A2 9/2008
 WO WO-2009/046220 A2 4/2009
 WO WO-2009/086558 A1 7/2009
 WO WO-2009/127060 A1 10/2009
 WO WO-2009/127230 A1 10/2009
 WO WO-2010/037408 A1 4/2010
 WO WO-2010/042877 A1 4/2010
 WO WO-2010/045512 A2 4/2010
 WO WO-2010/053572 A2 5/2010
 WO WO-2010/054401 A1 5/2010
 WO WO-2010/054405 A1 5/2010
 WO WO-2010/056403 A1 5/2010
 WO WO-2010/088537 A2 8/2010
 WO WO-2010/099387 A1 9/2010
 WO WO-2010/114789 A1 10/2010
 WO WO-2010/119256 A1 10/2010
 WO WO-2010/129709 A1 11/2010
 WO WO-2010/144740 A1 12/2010
 WO WO-2010/147992 A1 12/2010
 WO WO-2010/148013 A2 12/2010
 WO WO-2011/012316 A2 2/2011
 WO WO-2011/012746 A2 2/2011
 WO WO-2011/039144 A1 4/2011
 WO WO-2011/068810 A1 6/2011
 WO WO-2011/075656 A1 6/2011
 WO WO-2011/141705 A1 11/2011
 WO WO-2012/019168 A2 2/2012
 WO WO-2012/019630 A1 2/2012
 WO WO-2012/019780 A1 2/2012
 WO WO-2012/027675 A2 3/2012
 WO WO-2012/045075 A1 4/2012
 WO WO-2012/045082 A2 4/2012
 WO WO-2012/075040 A2 6/2012

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO WO-2012/133737 A1 10/2012
 WO WO-2012/135025 A2 10/2012
 WO WO-2012/135805 A2 10/2012
 WO WO-2012/170889 A1 12/2012
 WO WO-2012/170930 A1 12/2012
 WO WO-2013/039857 A1 3/2013
 WO WO-2013/039861 A2 3/2013
 WO WO-2013/063468 A1 5/2013
 WO WO-2013/090186 6/2013
 WO WO-2013/101690 A1 7/2013
 WO WO-2013/102203 A1 7/2013
 WO WO-2013/126803 A1 8/2013
 WO WO-2013/130161 9/2013
 WO WO-2013/149140 A1 10/2013
 WO WO-2013/149141 A1 10/2013
 WO WO-2013/151663 A1 10/2013
 WO WO-2013/151664 A1 10/2013
 WO WO-2013/151666 A2 10/2013
 WO WO-2013/151667 A1 10/2013
 WO WO-2013/151668 A2 10/2013
 WO WO-2013/151670 A2 10/2013
 WO WO-2013/151671 A1 10/2013
 WO WO-2013/151672 A2 10/2013
 WO WO-2013/151736 A2 10/2013
 WO WO-2013/182683 A1 12/2013
 WO WO-2013/185067 A1 12/2013
 WO WO-2013/185069 A1 12/2013
 WO WO-2014/028487 A1 2/2014
 WO WO-2014/089486 A1 6/2014
 WO WO-2014/113089 A2 7/2014
 WO WO-2014/144039 A1 9/2014
 WO WO-2014/144196 A1 9/2014
 WO WO-2014/144711 A1 9/2014
 WO WO-2014/144767 A1 9/2014
 WO WO-2014/152027 A1 9/2014
 WO WO-2014/152030 A1 9/2014
 WO WO-2014/152031 A1 9/2014
 WO WO-2014/152211 A1 9/2014
 WO WO-2014/152513 A1 9/2014
 WO WO-2014/152540 A1 9/2014
 WO WO-2014/152659 A1 9/2014
 WO WO-2014/152673 A1 9/2014
 WO WO-2014/152774 A1 9/2014
 WO WO-2014/152940 A1 9/2014
 WO WO-2014/152966 A1 9/2014
 WO WO-2014/153052 A2 9/2014
 WO WO-2014/158795 A1 10/2014
 WO WO-2014/159813 A1 10/2014
 WO WO-2014/17956-2 A1 11/2014
 WO WO-2014/210356 A1 12/2014
 WO WO-2015/006747 A2 1/2015
 WO WO-2015/011633 A1 1/2015
 WO WO-2015/048744 A2 4/2015
 WO WO-2015/051169 A2 4/2015
 WO WO-2015/051173 A2 4/2015
 WO WO-2015/058069 A1 4/2015
 WO WO-2015/085318 6/2015
 WO WO-2015/089511 6/2015
 WO WO-2016/054421 4/2016
 WO WO-2016/071857 A1 5/2016
 WO WO-2016/077123 A1 5/2016
 WO WO-2016/077125 5/2016
 WO WO-2016/077125 A1 5/2016
 WO WO-2016/118724 7/2016
 WO WO-2016/118725 7/2016
 WO WO-2016/154127 9/2016
 WO WO-2016/164762 10/2016
 WO WO-2016/183366 A2 11/2016
 WO WO-2016/197132 A1 12/2016
 WO WO-2016/197133 A1 12/2016
 WO WO-2016/201377 A1 12/2016
 WO WO-2017/019891 A2 2/2017
 WO WO-2017/049074 A1 3/2017
 WO WO-2017/049286 A1 3/2017
 WO WO-2017049275 A2 3/2017

WO WO-2017102010 A1 6/2017
 WO WO-2017103088 A1 6/2017
 WO WO-2017108087 A1 6/2017
 WO WO-2017109134 A1 6/2017
 WO WO-2017109161 A1 6/2017
 WO WO-2017117528 A1 7/2017
 WO WO-2017117530 A1 7/2017
 WO WO-2019207060 A1 10/2019

OTHER PUBLICATIONS

U.S. Appl. No. 61/494,714.
 U.S. Appl. No. 61/494,745.
 U.S. Appl. No. 61/494,881.
 U.S. Appl. No. 61/494,882.
 Adami, R.C. et al., An amino acid-based amphoteric liposomal delivery system for systemic administration of siRNA. *Molecular Therapy* 19(6):1141-1151 (2011).
 Akinc, A. et al., A combinatorial library of lipid-like materials for delivery of RNAi therapeutics. *Nature Biotechnology* 26(5):561-569 (2008).
 Akinc, A. et al., Development of lipidoid-siRNA formulations for systemic delivery to the liver. *Molecular Therapy* 17(5):872-879 (2009).
 Alton, E.W.F.W. et al., Cationic Lipid-Mediated CFTR Gene Transfer to the Lungs and Nose of Patients with Cystic Fibrosis: a Double-Blind Placebo-Controlled Trial, *Lancet*, 353:947-954 (1999).
 Anderson, D.G. et al., Structure/property studies of polymeric gene delivery using a library of poly(beta-amino esters). *Molecular Therapy* 11(3):426-434 (2005).
 Anderson, D.M. et al., Stability of mRNA/Cationic Lipid Lipoplexes in Human and Rat Cerebrospinal Fluid: Methods and Evidence for Nonviral mRNA Gene Delivery to the Central Nervous System, *Human Gene Therapy*, 14:191-202 (2003).
 Anderson, J. Biological Responses to Materials. *Annual Review of Materials Research* 31:81-110 (2001).
 Anderson, W. French, Human gene therapy, *Nature*, 392, 25-30 (1998).
 Andries, O. et al., Comparison of the Gene Transfer Efficiency of mRNA/GL67 and pDNA/GL67 Complexes in Respiratory Cells, *Mol. Pharmaceut.*, 9: 2136-2145 (2012).
 Auffray, C. et al., Purification of Mouse Immunoglobulin Heavy-Chain Messenger RNAs from Total Myeloma Tumor RNA, *European Journal of Biochemistry*, 107(2):303-314 (1980).
 Author Unknown, Blood Proteins, published by Wikipedia, San Francisco, CA, 2 pages, <http://en.wikipedia.org/wiki/Blood_proteins> downloaded May 17, 2015.
 Bahlke, M. A. et al., Progress towards in vivo use of siRNAs, *Molecular Therapy*, 13:644-670 (2006).
 Bajaj, A. et al., Synthesis and gene transfection efficacies of PEI-cholesterol-based lipopolymers. *Bioconjugate Chemistry* 19(8):1640-516511 (2008).
 Barreau, C. et al., Liposome-mediated RNA transfection should be used with caution, *RNA*, 12:1790-1793 (2006).
 Behike, et al., Progress towards in vivo Use of siRNAs, *Molecular Therapy*, 13:644-70 (2006).
 Behr, J. et al., Efficient Gene Transfer into Mammalian Primary Endocrine Cells with Lipo Polyamine-Coated DNA, *Proc. Nat.'l Acad. Sci.*, 86: 6982-6986 (1989).
 Bennett, J. Immune response following intraocular delivery of recombinant viral vectors, *Gene Therapy*, 10: 977-982 (2003).
 Bhaduri, S. et al., Procedure for the preparation of milligram quantities of adenovirus messenger ribonucleic acid, *J. Virol.*, 10(6): 1126-1129 (1972).
 Bloomfield, V.A., Quasi-Elastic Light Scattering Applications in Biochemistry and Biology, *Ann. Rev. Biophys. Bioeng.* 10:421-450 (1981).
 Boussif, O. et al., A versatile vector for gene and oligonucleotide transfer into cells in culture and in vivo: polyethylenimine. *Proceedings of the National Academy of Sciences of the USA*. 92(16):7297-7301 (1995).

(56)

References Cited

OTHER PUBLICATIONS

- Braun, C.S. et al., Structure/function relationships of polyamidoamine/DNA dendrimers as gene delivery vehicles. *Journal of Pharmaceutical Sciences* 94(2):423-436 (2005).
- Breunig, M. et al., Breaking up the correlation between efficacy and toxicity for nonviral gene delivery. *Proceedings of the National Academy of Sciences of the U S A.* 104(36):14454-14459 (2007).
- Breunig, M. et al., Mechanistic investigation of poly(ethylene imine)-based siRNA delivery: disulfide bonds boost intracellular release of the cargo. *Journal of Controlled Release* 130(1):57-63 (2008).
- Brey, D.M. et al., Controlling poly(beta-amino ester) network properties through macromer branching. *Acta Biomaterialia* 4(2):207-217 (2008).
- Brey, D.M. et al., Influence of macromer molecular weight and chemistry on poly(beta-amino ester) network properties and initial cell interactions. *Journal of Biomedical Materials Research Part A* 85(3):731-741 (2007).
- Budker, V. et al., Protein/Amphipathic Polyamine Complexes Enable Highly Efficient Transfection with Minimal Toxicity, *BioTechniques*, 23: 139-147 (1997).
- Burger, G. et al., Sequencing complete mitochondrial and plastid genomes, *Nature Protocols*, 2: 603-614 (2007).
- Burnett, J.C. et al., Current progress of siRNA/shRNA therapeutics in clinical trials. *Biotechnology Journal* 6(9):1130-1146 (2011).
- Burney et al., "Gene therapy for treatment of cystic fibrosis", *The Application of Clinical Genetics*, vol. 5, pp. 29-36, (2012).
- Byk, G. et al., Synthesis, activity, and structure-activity relationship studies of novel cationic lipids for DNA transfer. *Journal of Medical Chemistry* 41(2):224-235 (1998).
- Caplen, N.J. et al., In vitro liposome-mediated DNA transfection of epithelial cell lines using the cationic liposome DC-Chol/DOPE, *Gene Therapy*, 2:603-613 (1995).
- Cassiman, D. Gene transfer for inborn errors of metabolism of the liver: the clinical perspective, *Current Pharmaceutical Design*, 17(24):2550-2557 (2011).
- Castanotto, D. et al., The promises and pitfalls of RNA-interference-based therapeutics. *Nature* 457(7228):426-433 (2009).
- Chakraborty, C. Potentiality of Small Interfering RNAs (siRNA) as Recent Therapeutic Targets for Gene-Silencing. *Current Drug Targets* 8(3):469-82 (2007).
- Chandler, R. et al., Liver-directed adeno-associated virus serotype 8 gene transfer rescues a lethal murine model of citrullinemia type 1, *Gene Therapy*, 20:1188-1191 (2013).
- Chau, Y. et al., Investigation of targeting mechanism of new dextran-peptide-methotrexate conjugates using biodistribution study in matrix-metalloproteinase-overexpressing tumor xenograft model, *J. Pharm. Sci.*, 95(3): 542-551 (2006).
- Chen, D. et al., Rapid discovery of potent siRNA-containing lipid nanoparticles enabled by controlled microfluidic formulation. *Journal of the American Chemical Society* 134(16):6948-6951 (2012).
- Chen, Y. and Huang, L., Tumor-targeted delivery of siRNA by non-viral vector: safe and effective cancer therapy. *Expert Opinion on Drug Delivery* 5(12):1301-1311 (2008).
- Chiou, H.C. et al., Enhanced resistance to nuclease degradation of nucleic acids complexed to; asialoglycoprotein-polylysine carriers, *Nucleic Acids Research*, 22(24):5439-46 (1994).
- Christensen, U.B. et al., Intercalating nucleic acids containing insertions of 1-O-(1-pyrenylmethyl)glycerol: stabilisation of dsDNA and discrimination of DNA over RNA, *Nucl. Acids. Res.*, 30(22): 4918-4925 (2002).
- Conese, M. et al., Gene and Cell Therapy for Cystic Fibrosis: From Bench to Bedside, *J. Cyst. Fibros.*, 10 Suppl 2:S114-s128 (2011).
- Cotten, M. et al., Receptor-mediated transport of DNA into eukaryotic cells. *Methods in Enzymology* 217 (H):618-644 (1993).
- Cowling, V.H., Regulation of mRNA cap methylation, *Biochemical Journal*, 425:295-302 (2010).
- Creusat, G. et al., Proton sponge trick for pH-sensitive disassembly of polyethylenimine-based siRNA delivery systems. *Bioconjugate Chemistry* 21(5):994-1002 (2010).
- Crooke, et al., *Liposomal Formulations for Nucleic Acid Delivery, Antisense Drug Technology: Principles, Strategies, and Applications, Second Edition, Ch. 9, 237-270 (2007).*
- Crooke, S.T. Molecular mechanisms of action of antisense drugs. *Biochimica et Biophysica Acta* 1489(1):31-44. Review (1999).
- Crystal, R.G. Transfer of genes to humans: early lessons and obstacles to success. *Science* 270(5235):404-410. Review (1995).
- Damen, M. et al., Delivery of DNA and siRNA by novel gemini-like amphiphilic peptides. *Journal of Controlled Release* 145(1):33-39 (2010).
- Dande, P. et al., Improving RNA interference in mammalian cells by 4'-thio-modified small interfering RNA (siRNA): effect on siRNA activity and nuclease stability when used in combination with 2'-O-alkyl modifications, *Journal of Medicinal Chemistry*, 49(5):1624-1634 (2006).
- Dassa et al., Expression of the alternative oxidase complements cytochrome c oxidase deficiency in human cells, *EMBO Mol. Med.*, 1(1): 30-36 (2009).
- Davis, M. E., The first targeted delivery of siRNA in humans via a self-assembling, cyclodextrin polymer-based nanoparticle: from concept to clinic. *Molecular Pharmaceutics* 6(3):659-668 (2009).
- Davis, M.E. et al., Evidence of RNAi in humans from systemically administered siRNA via targeted nanoparticles. *Nature* 464(7291):1067-1070 (2010).
- Debus, H. et al., Delivery of Messenger RNA Using Poly(ethylene imine)-poly(ethylene glycol)-Copolymer Blends for Polyplex Formation: Biophysical Characterization and In Vitro Transfection Properties, *J. Control. Rel.*, 148:334-343 (2010).
- Decher, G. Fuzzy Nanoassemblies: Toward Layered Polymeric Multicomposites. *Science* 277: 1232-1237 (1997).
- Demeshkina, N. et al., Interactions of the ribosome with mRNA and tRNA, *Current Opinion in Structural Biology*, 20(3):325-332 (2010).
- Denardo, S.J. et al., Enhanced Therapeutic Index of Radioimmunotherapy (RIT) in Prostate Cancer Patients Comparison of Radiation Dosimetry for 1,4,7,10-Tetraazacyclododecane-N,N',N'',N'''-Tetraacetic Acid (DOTA)-Peptide versus 2IT-DOTA Monoclonal Antibody Linkage for RIT1, *Clin. Cancer Res.*, 9: 3665s (2003).
- Dern, R.J. et al., Toxicity studies of pyrimethamine (daraprim). *The American Journal of Tropical Medicine and Hygiene* 4(2):217-220 (1955).
- Deshmukh, H. M and Huang, L., Liposome and polylysine mediated gene therapy. *New Journal of Chemistry* 21:113-124 (1997).
- Discher, B.M. et al., Polymersomes: tough vesicles made from diblock copolymers. *Science* 284(5417):1143-1146 (1999).
- Discher, D.E. and Eisenberg, A., Polymer vesicles. *Science* 297(5583):967-973. Review (2002).
- Dokka et al., "Oxygen Radical-Mediated Pulmonary Toxicity Induced by Some Cationic Liposomes", *Pharmaceutical Research, Pharmaceutical Research*, 17(5): 521-525 (2000).
- Dong, Y. et al., Lipopeptide nanoparticles for potent and selective siRNA delivery in rodents and nonhuman primates, *Proceedings of the National Academy of Sciences*, 111(11): 3955-3960 (2014).
- Driscoll, K.E. et al., Intratracheal instillation as an exposure technique for the evaluation of respiratory tract toxicity: uses and limitations, *Toxicol. Sci.*, 55(1): 24-35 (2000).
- Drummond, D.C. et al., Optimizing Liposomes for Delivery of Chemotherapeutic Agents to Solid Tumors, *Pharmacological Reviews*, 51(4): 691-743 (1999).
- Dwarki, V. et al., Cationic liposome-mediated RNA transfection, *Methods in Enzymology*, 217:644-654 (1993).
- Elbashir, S.M. et al., RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes & Development* 15: 188-200 (2001).
- Elton, C., *The Next Next Big Thing*, Boston Magazine, pp. 106-118 (Mar. 2013).
- Emlen, W. et al., Effect of DNA size and strandedness on the in vivo clearance and organ localization of DNA, *Clinical & Experimental Immunology*, 56:185-192 (1984).
- Eon-Duval, A. et al., Removal of RNA impurities by tangential flow filtration in an RNase-free plasmid DNA purification process, *Analytical Biochemistry*, 316(1):66-73 (2003).
- Ernst, N. et al., Interaction of Liposomal and Polycationic Transfection Complexes with Pulmonary Surfactant, *J. Gene. Med.*, 1:331-340 (1999).

(56)

References Cited

OTHER PUBLICATIONS

- Estimated Number of Animal and Plant Species on Earth, <http://www.factmonster.com/ipka/A0934288.html>, 2000-2014, 3 pages, (Retrieved Aug. 2, 2014).
- Ewert, K. et al., Cationic lipid-DNA complexes for gene therapy: understanding the relationship between complex structure and gene delivery pathways at the molecular level. *Current Medicinal Chemistry* 11(2): 133-149 (2004).
- Fath, S. et al., Multiparameter RNA and Codon Optimization: A Standardized Tool to Assess and Enhance Autologous Mammalian Gene Expression, *PLoS One*, 6(3):e17596 (14 pages) 2011.
- Fechter, P. et al., Recognition of mRNA cap structures by viral and cellular proteins, *Journal of General Virology*, 86:1239-1249 (2005).
- Feigner, P.L. et al., Lipofection: A Highly Efficient, Lipid-Mediated DNA-Transfection Procedure, *Proc. Natl. Acad.*, 84:7413-7417 (1987).
- Felgner, P.L. and Ringold, G.M., Cationic liposome-mediated transfection, *Nature*, 337(6205):387-388 (1989).
- Fenske, D.B. and Cullis, P., Liposomal nanomedicines. *Expert Opinion on Drug Delivery* 5(1):25-44 (2008).
- Fenske, David B. et al., Entrapment of Small Molecules and Nucleic Acid-Based Drugs in Liposomes, *Methods in Enzymology*, 2005, pp. 7-40, vol. 391.
- Fernandez, V. et al., Cross Flow Filtration of RNA Extracts by Hollow Fiber Membrane, *Acta Biotechnologica*, 12(1):49-56 (1992).
- Ferruti, P.F. and Barbucci, R., Linear amino polymers: Synthesis, protonation and complex formation. *Advances in Polymer Science* 58:55-92 (1984).
- Ferruti, P.F. et al., A novel modification of poly(L-lysine) leading to a soluble cationic polymer with reduced toxicity and with potential as a transfection agent. *Macromolecular Chemistry and Physics* 199:2565-2575 (1998).
- Fire, A. et al., Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391(6669):806-811 (1998).
- Fischer, D. et al., Effect of poly(ethylene imine) molecular weight and pegylation on organ distribution and pharmacokinetics; of polyplexes with oligodeoxynucleotides in mice, *Drug Metabolism and Disposition*, 32(9):983-92 (2004).
- Fumoto et al., Targeted Gene Delivery: Importance of Administration Routes, *Novel Gene Therapy Approaches*, 3-31 (2013).
- Furgeson, D.Y. et al., Modified linear polyethylenimine-cholesterol conjugates for DNA complexation. *Bioconjugate Chemistry* 14(4):840-847 (2003).
- Furgeson, D.Y. et al., Novel water insoluble lipoparticulates for gene delivery. *Pharmaceutical Research* 19(4): 382-390 (2002).
- Galipon, J. et al., Stress-induced lncRNAs evade nuclear degradation and enter the translational machinery, *Genes to Cells*, 18(5):353-368 (2013).
- Gao, X. et al., A novel cationic liposome reagent for efficient transfection of mammalian cells, *Biochemical and Biophysical Research Communications*, 179(1):280-285 (1991).
- Garbuzenko, O.B. et al., Intratracheal Versus Intravenous Liposomal Delivery of siRNA, Antisense Oligonucleotides and Anticancer Drug, *Pharmaceutical Research*, 26(2):382-394 (2009).
- Geraerts, M. et al., Upscaling of lentiviral vector production by tangential flow filtration, *Journal of Gene Medicine*, 7(10):1299-1310 (2005).
- Godbey, W.T. et al., Size matters: molecular weight affects the efficiency of poly(ethylenimine) as a gene delivery vehicle. *Journal of Biomedical Materials Research* 45(3):268-275 (1998).
- Gonzalez, H. et al., New class of polymers for the delivery of macromolecular therapeutics. *Bioconjugate Chemistry* 10(6):1068-1074 (1999).
- Gonzalez-Aseguinolaza, G. et al., Gene therapy of liver diseases: A 2011 perspective, *Clinics and Research in Hepatology and Gastroenterology*, 35(11):699-708 (2011).
- Gordon, N. Ornithine transcarbamylase deficiency: a urea cycle defect, *European Journal of Paediatric Neurology*, 7:115-121 (2003).
- Grayson, A.C.R. et al., Biophysical and structural characterization of polyethylenimine-mediated siRNA delivery in vitro. *Pharmaceutical Research* 23(8): 1868-1876 (2006).
- Grudzien, E. et al., Novel cap analogs for in vitro synthesis of mRNAs with high translational efficiency, *RNA Biology*, 10(9):1479-1487 (2004).
- Grudzien, E. et al., Novel cap analogs for in vitro synthesis of mRNAs with high translational efficiency, Cold Spring Harbor Laboratory Press, 10(9):1479-1487 (2004).
- Grunlan, M.A. et al., Synthesis of 1,9-bis[glycidylxypropyl]penta(1'H, 1'H, 2'H, 2'H-perfluoroalkylmethylsiloxane)s and copolymerization with piperazine. *Polymer* 45:2517-2523 (2004).
- Gupta, U. et al., A review of in vitro-in vivo investigations on dendrimers: the novel nanoscopic drug carriers. *Nanomedicine: Nanotechnology, Biology, and Medicine* 2(2):66-73 (2006).
- Gust, T.C. et al., RNA-containing adenovirus/polyethylenimine transfer complexes effectively transduce dendritic cells and induce antigen-specific T cell responses, *The Journal of Gene Medicine*, 6(4): 464-470 (2004).
- Guttman, M. et al., Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals, *Nature*, 458:223-227 (2009).
- Haensler, J. and Szoka, F., Polyamidoamine cascade polymers mediate efficient transfection of cells in culture. *Bioconjugate Chemistry* 4(5):372-379 (1993).
- Harada-Shiba, M. et al., Polyion complex micelles as vectors in gene therapy pharmacokinetics and in vivo; gene transfer, *Gene Therapy*, 9(6):407-14 (2002).
- Haskins, Gene Therapy for Lysosomal Storage Disorders (LSDs) in Large Animal Models, *ILAR J.*, 50(2): 112-121 (2009).
- Hata, A. et al., Isolation and Characterization of the Human Ornithine Transcarbamylase Gene: Structure of the 5'-End Region, *Journal of Biochemistry*, 100:717-725 (1986).
- Hattatsu, Nou To, *Brain and Development*, 2007, pp. 87-92, vol. 39, No. 2.
- Hecker, J. et al., Advances in Self-Limited Gene Expression of Protective Intracellular Proteins In-Vivo in Rat Brain Using mRNA / Cationic Lipid Complexes, *Anesthesia and Analgesia*, 86(2S):346S (1994).
- Heidenreich, O. et al., High Activity and Stability of Hammerhead Ribozymes Containing 2'-Modified Pyrimidine Nucleosides and Phosphorothioates, *The Journal of Biological Chemistry*, 269(3):2131-2138 (1994).
- Henkin et al., Inhaled Insulin—Intrapulmonary, intranasal, and other routes of administration: Mechanisms of action, *Nutrition*, 26: 33-39 (2010).
- Hess, et al., Vaccination with mRNAs encoding tumor-associated antigens and granulocyte-macrophage colony-stimulating factor efficiently primes CTL responses, but is insufficient to overcome tolerance to a model tumor/self antigen, *Cancer Immunology, Immunotherapy: CII.* (2006), 55(6): 672-83.
- Heyes, J. et al., Cationic Lipid Saturation Influences Intracellular Delivery of Encapsulated Nucleic Acids, *J. Controlled Release*, 107:276-287 (2005).
- Higman, M.A. et al., The mRNA (Guanine-7-)methyltransferase Domain of the Vaccinia Virus mRNA Capping Enzyme, *The Journal of Biological Chemistry*, 269(21):14974-14981 (1994).
- Hill, I.R.C. et al., In vitro cytotoxicity of poly(amidoamine)s: relevance to DNA delivery. *Biochimica et Biophysica Acta* 1427: 161-174 (1999).
- Hill, J.G. et al., Enantioselective Epoxidation of Allylic Alcohols: (2S,3S)-3-Propyloxiranemethanol. *Organic Syntheses Collection* 7: 461 (1990) and 63: 66 (1985) (8 pages).
- Hillery, A.M. et al., *Drug Delivery and Targeting for Pharmacists and Pharmaceutical Scientists*, Taylor and Francis (2005).
- Hoerr, I. et al., In Vivo Application of RNA Leads to Induction of Specific Cytotoxic T Lymphocytes and Antibodies, *European Journal of Immunology*, 30(1):1-7 (2000).
- Hofland, H.E.J et al., Formation of stable cationic lipid/DNA complexes for gene transfer. *Proceedings of the National Academy of Sciences of the USA* 93 (14): 7305-7309 (1996).
- Homo sapiens* galactosidase, alpha (GLA) mRNA, NCBI Reference Sequence NM_000169.1, Modification Date: Nov. 17, 2006.

(56)

References Cited

OTHER PUBLICATIONS

- Hope, M.J. et al., Cationic Lipids, Phosphatidylethanolamine and the Intracellular Delivery of Polymeric, Nucleic Acid-Based Drugs. *Molecular Membrane Technology* 15:1-14 (1998).
- Hope, M.J. et al., Reduction of Liposome Size and Preparation of Unilamellar Vesicles by Extrusion Techniques, In: *Liposome Technology*, 1:123-139 (1993).
- Hornung, V. et al., Quantitative expression of toll-like receptor 1-10 mRNA in cellular subsets of human peripheral blood mononuclear cells and sensitivity to CpG oligodeoxynucleotides. *The Journal of Immunology* 168: 4531-4537 (2002).
- Horwich, A.L. et al., Structure and Expression of a Complementary DNA for the Nuclear Coded Precursor of Human Mitochondrial Ornithine Transcarbamylase, *Science*, 224(4653):1068-1074 (1984).
- Horwich, A.L. et al., Targeting of Pre-Ornithine Transcarbamylase to Mitochondria: Definition of Critical Regions and Residues in the Leader Peptide, *Cell*, 44:451-459 (1986).
- Howard, K.A. Delivery of RNA interference therapeutics using polycation-based nanoparticles. *Advanced Drug Delivery Reviews* 61: 710-720 (2009).
- Huang, Z. et al., Thiocholesterol-based lipids for ordered assembly of bioresponsive gene carriers, *Molecular Therapy*, 11(3):409-417 (2005).
- Huttenhofer, A. and Noller, H., Footprinting mRNA-ribosome complexes with chemical probes, *The EMBO Journal*, 13(16):3892-3901 (1994).
- Immordino, et al., "Stealth liposomes: review of the basic science, rationale, and clinical applications, existing and potential", *International Journal of Nanomedicine*, 1(3):297-315 (2006).
- Incani, V. et al., Lipid and hydrophobic modification of cationic carriers on route to superior gene vectors. *Soft Matter* 6: 2124-2138 (2010).
- International Preliminary Report on Patentability for PCT/US2010/058457, 12 pages (dated Jun. 14, 2012).
- International Search Report for PCT/US2010/58457, 4 pages (dated May 6, 2011).
- International Search Report for PCT/US2011/62459, 3 pages (dated Apr. 11, 2012).
- International Search Report for PCT/US2012/041663, 4 pages (dated Oct. 8, 2012).
- International Search Report for PCT/US2012/41724, 5 pages (dated Oct. 25, 2012).
- International Search Report for PCT/US2013/034602, 2 pages (dated Jun. 17, 2013).
- International Search Report for PCT/US2013/034604, 4 pages (dated Jun. 17, 2013).
- International Search Report for PCT/US2013/044769, 4 pages (dated Nov. 12, 2013).
- International Search Report for PCT/US2013/044771, 6 pages (dated Nov. 1, 2013).
- International Search Report for PCT/US2013/073672, 6 pages (dated Mar. 3, 2014).
- International Search Report for PCT/US2014/027422, 5 pages (dated Jul. 31, 2014).
- International Search Report for PCT/US2014/027585, 3 pages (dated Jul. 14, 2014).
- International Search Report for PCT/US2014/027587, 6 pages (dated Jul. 24, 2014).
- International Search Report for PCT/US2014/027602, 6 pages (dated Jul. 28, 2014).
- International Search Report for PCT/US2014/027717, 5 pages (dated Jul. 16, 2014).
- International Search Report for PCT/US2014/028330, 5 pages (dated Jul. 22, 2014).
- International Search Report for PCT/US2014/028441, 6 pages (dated Jul. 22, 2014).
- International Search Report for PCT/US2014/028498, 5 pages (dated Jul. 28, 2014).
- International Search Report for PCT/US2014/028849, 6 pages (dated Jul. 17, 2015).
- International Search Report for PCT/US2014/061786, 6 pages (dated Feb. 6, 2015).
- International Search Report for PCT/US2014/061793, 4 pages (dated Feb. 6, 2015).
- International Search Report for PCT/US2014/061830, 5 pages (dated Feb. 4, 2015).
- International Search Report for PCT/US2014/061841, 6 pages (dated Feb. 24, 2015).
- International Search Report for PCT/US2015/021403, 4 pages (dated Jun. 15, 2015).
- International Search Report for PCT/US2015/039004, 4 pages (dated Oct. 6, 2015).
- International Search Report for PCT/US2015/27563, 5 pages (dated Sep. 18, 2015).
- Jakobsen, K. et al., Purification of mRNA Directly From Crude Plant Tissues in 15 Minutes Using Magnetic Oligo DT Microspheres, *Nucleic Acids Research*, 18(12):3669 (1990).
- Jeffs, L.B. et al., A scalable, extrusion-free method for efficient liposomal encapsulation of plasmid DNA, *Pharmacol. Res.*, 22(3): 362-372 (2005).
- Jemielity, J. et al., Novel "anti-reverse" cap analogs with superior translational properties, Cold Spring Harbor Laboratory Press, 9(9):1108-1122 (2003).
- Jia, S. et al., "Eradication of osteosarcoma lung metastases following intranasal interleukin-12 gene therapy using a nonviral polyethylenimine vector," *Cancer Gene Therapy*, 9(3): 260-266 (2002).
- Jiang, G. et al., Hyaluronic acid-polyethyleneimine conjugate for target specific intracellular delivery of siRNA. *Biopolymers* 89 (7): 635-642 (2008).
- Jiang, M. et al., Electrochemically controlled release of lipid/DNA complexes: a new tool for synthetic gene delivery system. *Electrochemistry Communications* (6): 576-582 (2004).
- Jiang, S. and Cao, Z., Ultralow-fouling, functionalizable, and hydrolyzable zwitterionic materials and their derivatives for biological applications. *Advanced Materials* 22(9):920-932 (2010).
- Jolck, R.I. et al., Solid-phase synthesis of PEGylated lipopeptides using click chemistry. *Bioconjugate Chemistry* 21(5):807-810 (2010).
- Jon, S. et al., Degradable poly(amino alcohol esters) as potential DNA vectors with low cytotoxicity. *Biomacromolecules* 4(6):1759-1762 (2003).
- Jones, G. et al., Duplex- and Triplex-Forming Properties of 4'-Thio-Modified Oligodeoxynucleotides, *Bioorganic & Medicinal Chemistry Letters*, 7(10):1275-1278 (1997).
- Journal of the Japanese Society of Internal Medicine*, 2009, pp. 875-882, vol. 98, No. 4.
- Kabanov, A.V. and Kabanov, V.A., DNA complexes with polycations for the delivery of genetic material into cells. *Bioconjugate Chemistry* 6(1): 7-20 (1995).
- Kamath, S. et al., Surface chemistry influences implant-mediated host tissue responses. *Journal of Biomedical Materials Research A* 86(3):617-626 (2007).
- Kariko, et al., *Molecular Therapy*, (2008), 16(11): 1833-40.
- Kariko, K. et al., In vivo protein expression from mRNA delivered into adult rat brain, *Journal of Neuroscience Methods*, 105:77-86 (2001).
- Kasuya, T. et al., In Vivo Delivery of Bionanocapsules Displaying *Phaseolus vulgaris* Agglutinin-L₄ Isolectin to Malignant Tumors Overexpressing N-Acetylglucosaminyltransferase V, *Human Gene Therapy*, 19:887-895 (2008).
- Kaur, N. et al., A delineation of diketopiperazine self-assembly processes: understanding the molecular events involved in Nepsilon-(fumaroyl)diketopiperazine of L-Lys (FDKP) interactions. *Molecular Pharmaceutics* 5(2):294-315 (2007).
- Kaur, T. et al., Addressing the Challenge: Current and Future Directions in Ovarian Cancer Therapy, *Current Gene Therapy*, 9: 434-458 (2009).
- Kiew, L.V. et al., Effect of antisense oligodeoxynucleotides for ICAM-1 on renal ischaemia-reperfusion injury in the anaesthetised rat, *The Journal of Physiology*, 557(3):981-989 (2004).
- Kim, S.H. et al., Comparative evaluation of target-specific GFP gene silencing efficiencies for antisense ODN, synthetic siRNA, and

(56)

References Cited

OTHER PUBLICATIONS

- siRNA plasmid complexed with PEI-PEG-FOL conjugate. *Bioconjugate Chemistry* 17(1): 241-244 (2006).
- Kim, T. et al., Synthesis of biodegradable cross-linked poly(beta-amino ester) for gene delivery and its modification, inducing enhanced transfection efficiency and stepwise degradation. *Bioconjugate Chemistry* 16(5):1140-1148 (2005).
- Klibanov, A. et al., Amphipathic polyethyleneglycols effectively prolong the circulation time of liposomes. *Federation of European Biochemical Societies* 268 (1): 235-237 (1990).
- Kober, L. et al., Optimized Signal Peptides for the Development of High Expressing CHO Cell Lines, *Biotechnol. Bioeng.*, 110:1164-1173 (2012).
- Kodama, K. et al., The Features and Shortcomings for Gene Delivery of Current Non-Viral Carriers, *Current Medicinal Chemistry*, 13: 2155-2161 (2006).
- Kore, A. and Charles, I., Synthesis and evaluation of 2'-O-allyl substituted dinucleotide cap analog for mRNA translation, *Bioorganics & Medicinal Chemistry*, 18:8061-8065 (2010).
- Kore, A. and Shanmugasundaram, M., Synthesis and biological evaluation of trimethyl-substituted cap analogs, *Bioorganic & Medicinal Chemistry*, 18:880-884 (2008).
- Kormann, M.S.D. et al., Expression of therapeutic proteins after delivery of chemically modified mRNA in mice, *Nature Biotechnology*, 29(2):154-157 (2011).
- Kozak, M. An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs, *Nucleic Acid Research*, 15(20):8125-8148 (1987).
- Krieg, P.A. et al., In vitro RNA synthesis with SP6 RNA polymerase, *Methods in Enzymology*, 155:397-415 (1987).
- Kubo, et al., Mismatch Repair Protein Deficiency Is a Risk Factor for Aberrant Expression of HLA Class I Molecules: A Putative "Adaptive Immune Escape" Phenomenon, *Anticancer Res.*, 37(3): 1289-1295 (2017).
- Kvasnica, M. et al., Platinum(II) complexes with steroidal esters of L-methionine and L-histidine: Synthesis, characterization and cytotoxic activity, *Bioorganic & Medicinal Chemistry*, 16:3704-3713 (2008).
- Lam, J.K.W et al., Pulmonary delivery of therapeutic siRNA, *Advanced Drug Delivery Reviews* (2011).
- Lasic, D.D. et al., Gelation of liposome interior: A novel method for drug encapsulation, *FEBS*, 312(2,3):255-258 (1992).
- Lasic, D.D. Novel applications of liposomes, *Trends in Biotechnology*, 16:307-321 (1998).
- Lee, S. et al., Stability and cellular uptake of polymerized siRNA (poly-siRNA)/polyethylenimine (PEI) complexes for efficient gene silencing. *Journal of Controlled Release* 141: 339-346 (2010).
- Li, L. et al., Preparation and Gene Delivery of Alkaline Amino Acids-Based Cationic Liposomes, *Archives of Pharmaceutical Research*, 31(7):924-931 (2008).
- Li, S. et al., In vivo gene transfer via intravenous administration of cationic lipid-protamine-DNA (LPD) complexes, *Gene Therapy*, 4:891-900 (1997).
- Li, S.D., et al., Nanoparticles Evading The Reticuloendothelial System: Role of The Supported Bilayer; *Biochim Biophys Acta.*, Oct. 2009; 1788(10): 2259-2266.
- Li, W. et al., Lipid-based Nanoparticles for Nucleic Acid Delivery, *Pharmaceutical Research*, 24(3):438-449 (2007).
- Li, Yingfu et al., Kinetics of RNA Degradation by Specific Base Catalysis of Transesterification Involving the 2'-Hydroxyl Group, *J. Am. Chem. Soc.*, Feb. 1999, pp. 5364-5372, vol. 121.
- Liebhauer, S.A. et al., Translation inhibition by an mRNA coding region secondary structure is determined by its proximity to the AUG initiation codon, *Journal of Molecular Biology*, 226(3):609-621 (1992).
- Lim, Y. et al., A self-destroying polycationic polymer: biodegradable poly(4-hydroxy-L-proline ester). *Journal of American Chemical Society* 121: 5633-5639 (1999).
- Lindgren, V. et al., Human Ornithine Transcarbamylase Locus Mapped to Band Xp21.1 Near the Duchenne Muscular Dystrophy Locus, *Science*, 226(2675):698-700 (1984).
- Liu, X. et al., COStar: a D-star Lite-based Dynamic Search Algorithm for Codon Optimization, *Journal of Theoretical Biology*, 344:19-30 (2014).
- Liu, Y. et al., Designer Lipids Advance Systemic siRNA Delivery, *Molecular Therapy*, 18(4):669-670 (2010).
- Liu, Y. et al., Factors influencing the efficiency of cationic liposome-mediated intravenous gene delivery, *Nature Biotechnology*, 15:167-173 (1997).
- Lo, K-M et al., High level expression and secretion of Fc-X fusion proteins in mammalian cells, *Protein Engineering*, 11(6):495-500 (1998).
- Lorenzi, et al., Intranasal vaccination with messenger RNA as a new approach in gene therapy: use against tuberculosis, *BMC Biotechnology*, (Oct. 20, 2010), 10:77: pp. 1-11.
- Love, K.T. et al., Lipid-like materials for low-dose, in vivo gene silencing. *Proceedings of the National Academy of Sciences of the USA* 107 (5): 1864-1869 (2010).
- Lu, D. et al., Optimization of methods to achieve mRNA-mediated transfection of tumor cells in vitro and in vivo employing cationic liposome vectors, *Cancer Gene Therapy*, 1(4):245-52 (1994).
- Lukyanov, A.N. and Torchilin, V.P., Micelles from lipid derivatives of water-soluble polymers as delivery systems for poorly soluble drugs. *Advanced Drug Delivery Reviews* 56: 1273-1289 (2004).
- Luo, D. and Saltzman, M., Synthetic DNA delivery systems. *Nature Biotechnology* 18: 33-37. Review (2000).
- Lynn, D.M. and Langer, R., Degradable Poly(beta-amino esters):? Synthesis, Characterization, and Self-Assembly with Plasmid DNA. *Journal of American Chemical Society* 122(44): 10761-10768 (2000).
- Lynn, D.M. et al., Accelerated discovery of synthetic transfection vectors: parallel synthesis and screening of a degradable polymer library. *Journal of American Chemical Society* 123(33): 8155-8156 (2001).
- Lynn, D.M. et al., pH-Responsive Polymer Microspheres: Rapid Release of Encapsulated Material within the Range of Intracellular pH. *Angewandte Chemie International Edition* 40(9): 1707-1710 (2001).
- Ma, M. et al., Development of Cationic Polymer Coatings to Regulate Foreign Body Responses. *Advanced Healthcare Materials* 23: H189-H194. Reviews (2011).
- MacLachlan, I., Lipid nanoparticle-mediated delivery of messenger RNA, 1st International mRNA Health Conference; Tubingen Germany, (Oct. 24, 2013) Retrieved from the Internet: URL: <http://files.shareholder.com/downloads/ABEA-50QJTB/2628241206x0x699789/47543d12-db34-4e6e-88a9-f3ae5d97b1d2/MacLachlan_mRNA_Conf_2013>.
- Maeda-Mamiya, R. et al., In vivo gene delivery by cationic tetraamino; fullerene. *Proceedings of National Academy of Sciences U S A*, 107(12):5339-44 (2010).
- Malone, R.W., et al., Cationic liposome-mediated RNA transfection, *PNAS*, 86:6077-6081 (1989).
- Mammal, <http://en.wikipedia.org/wiki/Mammal>, 2007, Pearson Education, NY, NY, Author unknown (Source: The international union for conservation of nature and natural resources), 2 pages, (Retrieved Aug. 2, 2014).
- Mansour, H.M. et al., Nanomedicine in pulmonary delivery, *International Journal of Nanomedicine*, 4:299-319 (2009).
- Margus, H. et al., Cell-penetrating peptides as versatile vehicles for oligonucleotide delivery. *Molecular Therapy* 20 (3): 525-533 (2012).
- Martell, A.E. and Chaberek, S., The Preparation and the Properties of Some N,N'-Disubstituted-ethylenediaminedipropionic Acids. *Journal of the American Chemical Society* 72: 5357-5361 (1950).
- Martinon, et al., Induction of virus-specific cytotoxic T lymphocytes in vivo by liposome-entrapped mRNA, *European Journal of Immunology*, (1993), 23(7): 1719-22.
- Mathiowitz, E. and Langer, R., Polyanhydride microspheres as drug carriers I. Hot-melt microencapsulation. *Journal of Controlled Release* 5: 13-22 (1987).
- Mathiowitz, E. et al., Novel microcapsules for delivery systems. *Reactive Polymers* 6: 275-283 (1987).
- Mathiowitz, E. et al., Polyanhydride microspheres as drug carriers II. Microencapsulation by solvent removal. *Journal of Applied Polymer Sciences* 35: 755-774 (1988).

(56)

References Cited

OTHER PUBLICATIONS

- McCracken, S. et al., 5'-Capping Enzymes are Targeted to Pre-mRNA by Binding to the Phosphorylated Carboxy-Terminal Domain of RNA Polymerase II, *Genes and Development*, 22(24):3306-3318 (1997).
- McIvor, R. S., Therapeutic Delivery of mRNA: The Medium is the Message, *Molecular Therapy*, 19(5):822-823 (2011).
- Melton, D.A. et al., Efficient in vitro synthesis of biologically active RNA and RNA hybridization probes from plasmids containing a bacteriophage SP6 promoter, *Nucleic Acids Research*, 12(18):7035-56 (1984).
- Mendelsohn, J.D. et al., Rational design of cytophilic and cytophobic polyelectrolyte multilayer thin films. *Biomacromolecules* 4(1): 96-106 (2003).
- Merkel, O.M. et al., Nonviral Pulmonary Delivery of siRNA, *Accounts of Chemical Research*, 10 pages (2011).
- Merten, O. et al., Large-Scale Manufacture and Characterization of a Lentiviral Vector Produced for Clinical Ex Vivo Gene Therapy Application, *Human Gene Therapy*, 22(3):343-356 (2011).
- Meyer, et al., "Cationic liposomes coated with polyethylene glycol as carriers for oligonucleotides," *Journal of Biological Chemistry*, 273: 15621-15627 (1998).
- Miller, A. Cationic Liposomes for Gene Therapy. *Angewandte Chemie International Edition* 37: 1768-1785 (1998).
- Monck, et al., Stabilized Plasmid-Lipid Particles: Pharmacokinetics and Plasmid Delivery to Distal Tumors following Intravenous Injection, *J. Drug Target*, 7(6): 439-452 (2000).
- Monia, B.P. et al., Evaluation of 2'-Modified Oligonucleotides Containing 2'-Deoxy Gaps as Antisense Inhibitors of Gene Expression, *The Journal of Biological Chemistry*, 268(19):14514-14522 (1993).
- Morrissey, D.V. et al., Potent and Persistent in vivo Anti-HBV Activity of Chemically Modified siRNAs, *Nat. Biotechnol.*, 23(8): 1003-1007 (2005).
- Narang, A.S. et al., Cationic lipids with increased DNA binding affinity for nonviral gene transfer in dividing and nondividing cells. *Bioconjugate Chemistry* 16(1): 156-168 (2005).
- Navarro, G. et al., Phospholipid-polyethylenimine conjugate-based micelle-like nanoparticles for siRNA delivery. *Drug Delivery and Translational Research* 1: 25-33 (2011).
- Neamark, A. et al., Aliphatic lipid substitution on 2 kDa polyethylenimine improves plasmid delivery and transgene expression. *Molecular Pharmaceutics* 6(6): 1798-1815 (2009).
- Ng, J. et al., LincRNAs join the pluripotency alliance, *Nature Genetics*, 42:1035-1036 (2010).
- Nguyen, D.N. et al., A novel high-throughput cell-based method for integrated quantification of type I interferons and in vitro screening of immunostimulatory RNA drug delivery. *Biotechnology and Bioengineering* 103(4): 664-675 (2009).
- Nguyen, D.N. et al., Drug delivery-mediated control of RNA immunostimulation. *Molecular Therapy* 17(9): 1555-1562 (2009).
- Nojima, T. et al., The Interaction between Cap-binding Complex and RNA Export Factor is Required for Intronless mRNA Export, *Journal of Biological Chemistry*, 282(21):15645-15651 (2007).
- Nori, A. et al., Tat-conjugated synthetic macromolecules facilitate cytoplasmic drug delivery to human ovarian carcinoma cells, *Bioconj. Chem.*, 14(1): 44-50 (2003).
- Okumura, K. et al., Bax mRNA therapy using cationic liposomes for human malignant melanoma, *The Journal of Gene Medicine*, 10:910-917 (2008).
- Optiz, Lennart et al., Impact of RNA Degradation on Gene Expression Profiling, *BMC Medical Genomics*, 2010, pp. 1-14, 3:36, <http://www.biooscientific.com/AIR-DNA-Fragmentation-Kit>.
- Otsuka, Y. et al., Identification of a Cytoplasmic Complex That Adds a Cap onto 5'-Monophosphate RNA, *Molecular and Cellular Biology*, 29(8):2155-2167 (2009).
- Ozawa, Keeiya, M.D., Ph.D., Gene Therapy Using AAV, 2007, pp. 47-56, Division of Hematology, Department of Medicine, Division of Geriatric Therapeutics, Center for Molecular Medicine, Jichi Medical University, 3311-1 Yakuhiji, Shimotsuke-shi, Tochigi 329-0198, Japan.
- Ozer, A., Alternative applications for drug delivery: nasal and pulmonary routes, *Nanomaterials and Nanosystems for Biomedical Applications*, M.R. Mozafari (ed.): 99-112 (2007).
- Painter, H. et al., Topical Delivery of mRNA to the Murine Lung and Nasal Epithelium, *Gene Medicine Group and the Medical Informatics Unit, Nuffield Department of Clinical Laboratory Sciences, University of Oxford*, 1 page.
- Painter, H. et al., Topical Delivery of mRNA to the Murine Lung and Nasal Epithelium, *Molecular Therapy*, 9:S187 (2004).
- Painter, H., An Investigation of mRNA as a Gene Transfer Agent, *Gene Medicine Research Group Nuffield Department of Clinical Laboratory Sciences and Merton College, University of Oxford*, 1-282 (2007).
- Painter, H., An Investigation of mRNA as a Gene Transfer Agent, *Oxford University GeneMedicine, Abstract Only*, 1 page (2007).
- Parrish, D.A. and Mathias, L.J., Five- and six-membered ring opening of pyroglutamic diketopiperazine. *Journal of Organic Chemistry* 67(6): 1820-1826 (2002).
- Pathak, et al., Nano-vectors for efficient liver specific gene transfer, *Int. J. Nanomedicine*, 3(1): 31-49 (2008).
- Patton, J., Market Trends in Pulmonary Therapies, *Trends and Opportunities*, VI: 372-377.
- Paulus, C. and Nevels, M., The Human Cytomegalovirus Major Immediate-Early Proteins as Antagonists of Intrinsic and Innate Antiviral Host Responses, *Viruses*, 1:760-779 (2009).
- Peppas, N.A. et al., Hydrogels in Biology and Medicine: From Molecular Principles to Bionanotechnology. *Advanced Materials* 18: 1345-1360 (2006).
- Philipp, A. et al., Hydrophobically modified oligoethylenimines as highly efficient transfection agents for siRNA delivery. *Bioconjugate Chemistry* 20(11): 2055-2061 (2009).
- Pons, M. et al., Liposomes obtained by the ethanol injection method, *Int. J. Pharm.*, 95: 51-56. (1993).
- Prata, C.A. et al., Lipophilic peptides for gene delivery. *Bioconjugate Chemistry* 19(2): 418-420 (2008).
- Probst, J. et al., Spontaneous cellular uptake of exogenous messenger RNA in vivo is nucleic acid-specific, saturable and ion dependent, *Gene Therapy*, 14:1175-1180 (2007).
- Promega, PolyAtract mRNA Isolation Systems, Instructions for Use of Products Z5200, Z5210, Z2300 and Z5310, *Technical Manual* (2012).
- Putnam, D. Polymers for gene delivery across length scales. *Nature Materials* 5: 439-451 (2006).
- Putnam, D. and Langer, R., Poly(4-hydroxy-L-proline ester): Low-Temperature Polycondensation and Plasmid DNA Complexation. *Macromolecules* 32(11): 3658-3662 (1999).
- Qiagen, Oligotex Handbook, Second Edition (2002).
- Rabinovich, P.M. et al., Synthetic Messenger RNA as a Tool for Gene Therapy, *Human Gene Therapy*, 17:1027-1035 (2006).
- Ramaswamy, S., et al., "Systemic delivery of factor IX messenger RNA for protein replacement therapy", *Proceedings of the National Academy of Sciences of the United States of America*, 114(10):E1941-E1950 (Mar. 7, 2017).
- Raney, et al. The effect of circulation lifetime and drug-to-lipid ratio of intravenously administered lipid nanoparticles on the biodistribution and immunostimulatory activity of encapsulated CpG-ODN, *J. Drug Target*, 16(7): 564-577 (2008).
- Raper, S.E. et al., Developing adenoviral-mediated in vivo gene therapy for ornithine transcarbamylase deficiency, *Journal of Inherited Metabolic Disease*, 21:119-137 (1998).
- Ratajczak, J. et al., Membrane-derived microvesicles: important and underappreciated mediators of cell-to-cell communication, *Leukemia*, 20:1487-1495 (2006).
- Ratner, B.D. and Bryant, S., Biomaterials: where we have been and where we are going. *Annual Review of Biomedical Engineering* 6: 41-75 (2004).
- Reddy, A. et al., The Effect of Labour and Placental Separation on the Shedding of Syncytiotrophoblast Microparticles, Cell-free DNA and mRNA in Normal Pregnancy and Pre-eclampsia, *Placenta*, 29:942-949 (2008).

(56)

References Cited

OTHER PUBLICATIONS

- Rejman, J. et al., Characterization and transfection properties of lipoplexes stabilized with novel exchangeable polyethylene glycol-lipid conjugates, *Biochimica et Biophysica Acta*, 1660:41-52 (2004).
- Remington: *The Science and Practice of Pharmacy*, 21st Edition, Philadelphia, PA. Lippincott Williams & Wilkins (2005).
- Riordan, Thomas G. MD., Formulations and Nebulizer Performance, *Respiratory Care*, Nov. 2002, pp. 1305-1312, vol. 47, No. 11.
- Robinson et al., "Lipid Nanoparticle-Delivered Chemically Modified mRNA Restores Chloride Secretion in Cystic Fibrosis", *Molecular Therapy*, 26(8): 1-13 (2018).
- Rosenecker, J. et al., Gene Therapy for Cystic Fibrosis Lung Disease: Current Status and Future Perspectives, *Curr. Opin. Mol. Ther.*, 8:439-445 (2006).
- Rosenecker, J. et al., Interaction of Bronchoalveolar Lavage Fluid with Polyplexes and Lipoplexes: Analysing the Role of Proteins and Glycoproteins, *J. Gene. Med.*, 5:49-60 (2003).
- Rowe, S.M. et al., Cystic Fibrosis, *New Engl. J. Med.* 352:1992-2001 (2005).
- Rudolph, C. et al., Aerosolized Nanogram Quantities of Plasmid DNA Mediate Highly Efficient Gene Delivery to Mouse Airway Epithelium, *Molecular Therapy*, 12(3): 493-501 (2005).
- Rudolph, C. et al., Methodological optimization of polyethylenimine (PEI)-based gene delivery to the lungs of mice via aerosol application, *Journal of Gene Medicine*, 7(1): 59-66 (2005).
- Ryng, S. et al., Synthesis and structure elucidation of 5-aminomethinimino-3-methyl-4-isoxazolecarboxylic acid phenylamides and their immunological activity. *Arch. Pharm. Pharm. Med. Chem* 330(11):319-26 (1997).
- Sahay, G. et al., Endocytosis of nanomedicines. *Journal of Controlled Release* 145: 182-195 (2010).
- Sakiyama-Elbert, S.E. and Hubbell, J.A., Functional Biomaterials: Design of Novel Biomaterials. *Annual Review of Materials Research* 31: 183-201 (2001).
- Sambrook, Joseph et al., Fragmentation of DNA by Nebulization, Commonly Used Techniques in Molecular Cloning, Appendix 8, in *Molecular Cloning*, 2001-2006, 5 Pages., vol. 3, 3rd edition (eds. Sambrook and Russell). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA.
- Scherphof, et al., Uptake of Liposomes by Rat and Mouse Hepatocytes and Kupffer Cells, *Biol. Cell*, 47: 47-58 (1983).
- Schnierle, B.S. et al., Cap-specific mRNA (nucleoside-O^{2'}-)-methyltransferase and poly(A) polymerase stimulatory activities of vaccinia virus are mediated by a single protein, *Proceedings of the National Academy of Sciences*, 89:2897-2901 (1992).
- Schreier, H., The new frontier: gene and oligonucleotide therapy, *Pharmaceutica Acta Helveticae*, 68(3):145-159 (1994).
- Semple, S.C. et al., Rational design of cationic lipids for siRNA delivery, *Nature Biotechnology*, 28(2): 172-176 (2010).
- Shchori E., Poly(secondary Amine)s from Diacrylates and Diamines. *Journal of Polymer Science* 21(6):413-15 (1983).
- Sherwood, R.F. Advanced drug delivery reviews: enzyme prodrug therapy, *Adv. Drug Del. Rev.*, 22: 269-288 (1996).
- Shimada, A. et al., Translocation Pathway of the Intratracheally Instilled Ultrafine Particles from the Lung into the Blood Circulation in the Mouse, *Toxicologic Pathology*, 34:949-957 (2006).
- Sieglwart, D.J. et al., Combinatorial synthesis of chemically diverse core-shell nanoparticles for intracellular delivery. *Proceedings of the National Academy of the Sciences of the USA* 108(32):12996-123001 (2011).
- Sigurgeirsson, Benjamin et al., Sequencing Degraded RNA Addressed by 3' Tag Counting, *PLOS One*, Mar. 2014, pp. 1-11, vol. 9, Issue 3, e91851.
- Smisterova, J. et al., Molecular Shape of the Cationic Lipid Controls the Structure of Cationic Lipid/Dioleoylphosphatidylethanolamine-DNA Complexes and the Efficiency of Gene Delivery, *The Journal of Biological Chemistry*, 276(50):47615-47622 (2001).
- Stern, L. et al., A novel antitumor prodrug platform designed to be cleaved by the endoprotease legumain, *Bioconj. Chem.*, 20: 500-510 (2009).
- Su, X. et al., Cytosolic Delivery Mediated Via Electrostatic Surface Binding of mRNA To Degradable Lipid-Coated Polymeric Nanoparticles, *Polymer Preprints*, 51(2):668-669 (2010).
- Su, X. et al., In Vitro and in Vivo mRNA Delivery Using Lipid-Enveloped pH-Responsive Polymer Nanoparticles, *Molecular Pharmaceutics*, 8(3):774-787 (2011).
- Suri, M. et al., *Genetics for Pediatricians*, Remedica Publishing, (2005).
- Szoka, F. and Papahadjopoulos, D., Comparative properties and methods of preparation of lipid vesicles (liposomes). *Annual Review of Biophysics Bioengineering* 9: 467-508 (1980).
- Tagawa, M. et al., Gene expression and active virus replication in the liver after injection of duck hepatitis B virus DNA into the peripheral vein of ducklings, *Journal of Hepatology*, 24:328-334 (1996).
- Takahashi, Y. et al., Development of safe and effective nonviral gene therapy by eliminating CpG motifs from plasmid DNA vector, *Frontiers in Bioscience*, S4: 133-141 (2012).
- Tan, S. et al., Engineering Nanocarriers for siRNA Delivery. *Small* 7(7): 841-856 (2011).
- Tang, F. and Hughes, J. et al., Introduction of a Disulfide Bond into a Cationic Lipid Enhances Transgene Expression of Plasmid DNA, *Biochemical and Biophysical Research Communications*, 242(1):141-145 (1998).
- Tang, M.X. et al., In vitro gene delivery by degraded polyamidoamine dendrimers. *Bioconjugate Chemistry* 7(6): 703-714 (1996).
- Tarcha, P.J. et al., Synthesis and characterization of chemically condensed oligoethylenimine containing beta-aminopropionamide linkages for siRNA delivery. *Biomaterials* 28: 3731-3740 (2007).
- Tavernier, G. et al., mRNA as gene therapeutic: How to control protein expression, *Journal of Controlled Release*, 150:238-247 (2011).
- Tcherepanova, I. et al., Ectopic expression of a truncated CD40L protein from synthetic post-transcriptionally capped RNA in dendritic cells induces high levels of IL-12 secretion, *BMC Molecular Biology*, 9(1):pp. 1-13 (2008).
- Thess, A., et al., "Sequence-engineered mRNA Without Chemical Nucleoside Modifications Enables an Effective Protein Therapy in Large Animals", *Molecular Therapy*, 23(9):1456-64 (Sep. 2015).
- Theus, S. and Liarakos, C., A Simple Assay for Determining the Capping Efficiencies of RNA Polymerases Used for In Vitro Transcription, *BioChromatography*, 9(5):610-614 (1990).
- Third Party Preissuance Submission Under 37 CFR § 1.290 (Oct. 25, 2013).
- Thomas, C. E. et al., Progress and problems with the use of viral vectors for gene therapy, *Nature Reviews/Genetics*, 4: 346-358 (2003).
- Thompson, P.E. et al., Antiamebic action of 5-chloro-7-diethylaminomethyl-8-quinolinol and of other substituted 8-quinolinols in vitro and in experimental animals. *American Journal of Tropical Medicine and Hygiene* 2(4): 224-248 (1955).
- Toki, B.E. et al., Protease-mediated fragmentation of p-amidobenzyl ethers: a new strategy for the activation of anticancer prodrugs, *J. Org. Chem.*, 67(6): 1866-1872 (2002).
- Tolmachev, Oleg E. et al., Methods of Transfection with Messenger RNA Gene Vectors, *Gene Therapy—Principles and Challenges*, 2015, pp. 1-55, Chapter 2. <http://dx.doi.org/10.5772/61688>.
- Tranchant, I. et al., Physicochemical optimisation of plasmid delivery by cationic lipids. *Journal of Gene Medicine* 6: S24-S35 (2004).
- Tsui, N.B. et al., Stability of endogenous and added RNA in blood specimens, serum, and plasma, *Clinical Chemistry*, 48(10):1647-53 (2002).
- Tsvetkov, D.E. et al., Neoglycoconjugates based on dendrimeric poly(aminoamides). *Russian Journal of Bioorganic Chemistry* 28(6): 470-486 (2002).
- Tuschl, T. et al., Targeted mRNA degradation by double-stranded RNA in vitro, *Genes and Development*, 13(24):3191-7 (1999).
- Urban-Klein, B. et al., RNAi-mediated gene-targeting through systemic application of polyethylenimine (PEI)-complexed siRNA in vivo. *Gene Therapy* 12(5): 461-466 (2005).

(56)

References Cited

OTHER PUBLICATIONS

- Van Balen, G.P. et al., Liposome/water lipophilicity: methods, information content, and pharmaceutical applications. *Medicinal Research Reviews* 24(3): 299-324 (2004).
- Van De Wetering, P. et al., Structure-activity relationships of water-soluble cationic methacrylate/methacrylamide polymers for nonviral gene delivery. *Bioconjugate Chemistry* 10(4): 589-597 (1999).
- Van Der Gun, B.T.F. et al., Serum insensitive, intranuclear protein delivery by the multipurpose cationic lipid Saint-2, *Journal of Controlled Release*, 123:228-238 (2007).
- Van Tendeloo, V.F.I. et al., mRNA-based gene transfer as a tool for gene and cell therapy, *Current Opinion in Molecular Therapeutics*, 9(5):423-431 (2007).
- Vandenbroucke, R.E. et al., Prolonged gene silencing in hepatoma cells and primary hepatocytes after small interfering RNA delivery with biodegradable poly(beta-amino esters). *Journal of Gene Medicine* 10: 783-794 (2008).
- Varambally, S. et al., Genomic Loss of microRNA-101 Leads to Overexpression of Histone Methyltransferase EZH2 in Cancer, *Science*, 322:1695-1699 (2008).
- Veronese, F.M. et al., PEG-doxorubicin conjugates: influence of polymer structure on drug release, in vitro cytotoxicity, biodistribution, and antitumor activity, *Bioconj. Chem.*, 16(4): 775-784 (2005).
- Viecelli, H. et al., Gene Therapy for Hepatic Diseases Using Non-Viral Minicircle-DNA Vector, *Journal of Inherited Metabolic Disease*, 35(1):S144 (2012).
- Viecelli, H. et al., Gene therapy for liver diseases using non-viral minicircle-DNA vector, *Human Gene Therapy*, 23(10):A145 (2012).
- Viecelli, H. et al., Gene therapy for liver diseases using non-viral minicircle-DNA vector, *Molecular Therapy*, 21(1):S136 (2013).
- Vomelova, I. et al., Methods of RNA Purification. All Ways (Should) Lead to Rome, *Folia Biologica*, 55(6):242-251 (2009).
- Von Harpe et al., Characterization of commercially available and synthesized polyethylenimines for gene delivery. *Journal of Controlled Release* 69(2):309-322 (2000).
- Walde, P. et al., Preparation of Vesicles (Liposomes). *Encyclopedia of Nanoscience and Nanotechnology*. Nalwa, ed. American Scientific Publishers, Los Angeles 9:43-79 (2004).
- Wang, H. et al., N-acetylgalactosamine functionalized mixed micellar nanoparticles for targeted delivery of siRNA to liver, *Journal of Controlled Release*, 166(2):106-114 (2013).
- Wang, Y. et al., Systemic delivery of modified mRNA encoding herpes simplex virus 1 thymidine kinase for targeted cancer gene therapy, *Molecular Therapy*, 21(2):358-367 (2013).
- Wang, Yaogeng et al., The Impact of Different Preservation Conditions and Freezing-Thawing Cycles on Quality of RNA, DNA and Proteins in Cancer Tissue, *Biopreservation and Biobanking*, 2015, pp. 335-347, vol. 13, No. 5.
- Webb, M. et al., Sphingomyeline-cholesterol liposomes significantly enhance the pharmacokinetic and therapeutic properties of vincristine in murine and human tumour models, *British Journal of Cancer*, 72(4):896-904 (1995).
- Werth, S. et al., A low molecular weight fraction of polyethylenimine (PEI) displays increased transfection efficiency of DNA and siRNA in fresh or lyophilized complexes. *Journal of Controlled Release* 112: 257-270 (2006).
- Wetzer, B. et al., Reducible cationic lipids for gene transfer, *Biochem. J.*, 356:747-756 (2001).
- White, J.E. et al., Poly(hydroxyaminoethers): A New Family of Epoxy-Based Thermoplastics. *Advanced Materials* 12(23): 1791-1800 (2000).
- White, J.E. et al., Step-growth polymerization of 10,11-epoxyundecanoic acid. Synthesis and properties of a new hydroxy-functionalized thermoplastic polyester. *Advanced Materials* 48: 3990-3998 (2007).
- Whitehead, K.A. et al., Knocking down barriers: advances in siRNA delivery. *Nature Reviews Drug Discovery* 8(2): 129-139 (2009).
- Wiehe, J.M. et al., mRNA-mediated gene delivery into human progenitor cells promotes highly efficient protein expression, *Journal of Cellular and Molecular Medicine*, 11(3):521-530 (2007).
- Williams, D. et al., A simple, highly efficient method for heterologous expression in mammalian primary neurons using cationic lipid-mediated mRNA transfection, *Frontiers in Neuroscience*, 4(181):1-20 (2010).
- Wilson, et al., Ex vivo gene therapy of familial hypercholesterolemia, *Hum. Gene Ther.*, 3(2): 179-222 (1992).
- Wisse, et al., The size of endothelial fenestrae in human liver sinusoids: implications for hepatocyte-directed gene transfer, *Gene Therapy*, 15: 1193-1199 (2008).
- Wolf, J.A. et al., Protein/Amphiphathic Polyamine Complexes Enable Highly Efficient Transfection with Minimal Toxicity, *BioTechniques*, 23:139-147 (1997).
- Woodle, et al., Versality in lipid compositions showing prolonged circulation with sterically stabilized liposomes, *Biochim. Biophys. Acta*, 1105(2): 193-200 (1992).
- Written Opinion for PCT/US2010/058457, 10 pages (dated May 6, 2011).
- Written Opinion for PCT/US2011/062459, 9 pages (dated Apr. 11, 2012).
- Written Opinion for PCT/US2012/041663, 7 pages (dated Oct. 8, 2012).
- Written Opinion for PCT/US2012/041724, 11 pages (dated Oct. 25, 2012).
- Written Opinion for PCT/US2013/034602, 4 pages (dated Jun. 17, 2013).
- Written Opinion for PCT/US2013/034604, 9 pages (dated Jun. 17, 2013).
- Written Opinion for PCT/US2013/044769, 8 pages (dated Nov. 12, 2013).
- Written Opinion for PCT/US2013/044771, 7 pages (dated Nov. 1, 2013).
- Written Opinion for PCT/US2013/073672, 7 pages (dated Mar. 3, 2014).
- Written Opinion for PCT/US2014/027422, 6 pages (dated Jul. 31, 2014).
- Written Opinion for PCT/US2014/027587, 5 pages (dated Jul. 24, 2014).
- Written Opinion for PCT/US2014/027602, 7 pages (dated Jul. 28, 2014).
- Written Opinion for PCT/US2014/027717, 5 pages (dated Jul. 16, 2014).
- Written Opinion for PCT/US2014/028330, 7 pages (dated Jul. 22, 2014).
- Written Opinion for PCT/US2014/028441, 6 pages (dated Jul. 22, 2014).
- Written Opinion for PCT/US2014/028498, 6 pages (dated Jul. 28, 2014).
- Written Opinion for PCT/US2014/028849, 7 pages (dated Jul. 17, 2015).
- Written Opinion for PCT/US2014/061786, 5 pages (dated Feb. 6, 2015).
- Written Opinion for PCT/US2014/061793, 4 pages (dated Feb. 6, 2015).
- Written Opinion for PCT/US2014/061830, 7 pages (dated Feb. 4, 2015).
- Written Opinion for PCT/US2014/061841, 8 pages (dated Feb. 24, 2015).
- Written Opinion for PCT/US2015/021403, 7 pages (dated Jun. 15, 2015).
- Written Opinion for PCT/US2015/027563, 12 pages (dated Sep. 18, 2015).
- Written Opinion for PCT/US2015/039004, 8 pages (dated Oct. 6, 2015).
- Wu, J. and Zern, M., Modification of liposomes for liver targeting, *Journal of Hepatology*, 24(6):757-763 (1996).
- Wu, J. et al., Cationic lipid polymerization as a novel approach for constructing new DNA delivery agents. *Bioconjugate Chemistry* 12(2): 251-257 (2001).
- Wurdinger, et al. (2008), A secreted luciferase for ex-vivo monitoring of vivo processes, *Nature Methods*, 5(2): 171-173.
- Yamamoto, A. et al., Current prospects for mRNA gene delivery, *European Journal of Pharmaceutics and Biopharmaceutics*, 71:484-489 (2009).

(56)

References Cited

OTHER PUBLICATIONS

Yamamoto, Y. et al., Important Role of the Proline Residue in the Signal Sequence that Directs the Secretion of Human Lysozyme in *Saccharomyces cerevisiae*, *Biochemistry*, 28:2728-2732 (1989).

Yang, Tzu-Hsueh et al., Determination of RNA degradation by Capillary Electrophoresis with Cyan Light-emitted Diode-induced Fluorescence, *Journal of Chromatography*, Mar. 2012, pp. 78-84, vol. 1239, Journal Homepage: www.elsevier.com/locate/chroma.

Yasuda et al., Fabry Disease: Novel [alpha]-Galactosidase A 3-terminal Mutations Result in Multiple Transcripts Due to Aberrant 3-End Formation, *American Journal of Human Genetics*, 73: 162-73 (2003).

Ye, X. et al., Nucleic Acids, Protein Synthesis, and Molecular Genetics: Prolonged Metabolic Correction in Adult Ornithine Transcarbamylase-deficient Mice with Adenoviral Vectors, *The Journal of Biological Chemistry*, 271:3639-3646 (1996).

Yokoe, H. et al., Spatial dynamics of GFP-tagged proteins investigated by local fluorescence enhancement, *Nature Biotechnology*, 14(10):1252-1256 (1996).

Yoneda et al., A cell-penetrating peptidic GRP78 ligand for tumor cell-specific prodrug therapy, *Bioorg. Med. Chem. Lett.*, 18(5): 1632-1636 (2008).

Yoshioka, Y. and Calvert, P., Epoxy-based Electroactive Polymer Gels. *Experimental Mechanics* 42(4): 404-408 (2002).

Zagridullin, P.H. et al., Monobasic amines. II. Cycloalkylation and hydroxyalkylation of cyclic and acyclic di- and polyamines. *Journal of Organic Chemistry* 26(1):184-88. Russian (1990).

Zaugg, H.E. et al., 3-Carboxy-2,5-piperazinedione and Derivatives. *Journal of American Chemical Society* 78(11):2626-2631 (1956).

Zauner, W. et al., Polylysine-based transfection systems utilizing receptor-mediated delivery. *Advanced Drug Delivery Reviews* 30(1-3):97-113(1998).

Zhang, L. et al., "Enhanced efficacy and limited systemic cytokine exposure with membrane-anchored interleukin-12 T-cell therapy in murine tumor models," *Journal for ImmunoTherapy of Cancer*, 8(1): article e000210 (2020).

Zintchenko, A. et al., Simple modifications of branched PEI lead to highly efficient siRNA carriers with low toxicity. *Bioconjugate Chemistry* 19(7):1448-1455 (2008).

Zou, S. et al., Lipid-mediated delivery of RNA is more efficient than delivery of DNA in non-dividing cells, *International Journal of Pharmaceutics*, 389(1-2):232-243 (2010).

5' CMV Sequence:

XCAGAUCCUGGAGAGCCCAUCCACGGUUGUUUGACUCCCAUAGAACACCCGGACCGAUCCAGCCUCCGGGGC
CGGAAACGGUGCAUUGAACGGGAUCCCGUGCCAAGAGUCUCUACCGUCCUUGACAG
(SEQ ID NO: 1)

FIG. 1

3' hGH Sequence:

CGGGUGGCAUCCUGUGACCCUCCAGUGCCUCCUGGCCUUGGAAGUUGCCACUCCAGUGGCCACCAGCCUUG
UCCUAUAAAUAUAGUUGCAUC (SEQ ID NO: 2)

FIG. 2

Human erythropoietin (EPO) mRNA:

AUGGGGUGCACGGAUGUUCUGCCUGGUGGCUUCCUGUCCUUGGCUCCUUGGCUCCUUGGCUCCAGUCC
UGGGGCCCCACCCAGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUUGGAGGCCAAGGAGGCCGAGAA
UAUCACGCGGCUUGUGAACACUCGACGCUUGAUGAGAAUACUCUCCAGACACCAAGUUAUUUCUUAU
GCCUGGAAGAGGAUGGAGGUCGGGCAGCCCGUAGAGUCUGGCGAGGCCUUGGCCUUGCGGAGGCUUCC
UGGGGGCCAGGCCUUGUUGUCAACUCUCCAGCCGUGGGAGCCUUGCAGCUGCAUGUGGAUAAAGCCGUCAG
UGGCCUUGCGAGCCUACCCACUCUGCUUGGGUCUGGAGCCAGAGGAGCCAUUCCUCCUCCAGAUUGGGCCU
CAGCUGUCCACUCCGACACAUAUCUGCUGACACUUCCGCAACUCUCCGAGUCUACUCCAAUUUCCUCCGGG
AAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGGACAGAUGA (SEQ ID NO: 3)

FIG. 3

Human alpha-galactosidase (GLA) mRNA:

AUGCAGCUGAGGAACCCAGAAACUACUUGGGCUGCGGUUGCCUUCGCUUCUGGCCUUCGUUUCUGGGACA
UCCUUGGGCUAGAGCACUGGACAUGGAUUGGCAAGGACGCCUACCAUGGGCUGGCUCACUGGGAGCGCUUCAU
GUGCAACCUGACUGCCAGGAGAGCCAGAUUCCUGCAUCAGUGAGAGCUUCUUCAGGAGAUUGGCAGAGCUC AUG
GUCUCAGAAGCCUGGAAGGAUGCAGGUUAGAUCCUCUGCAUUGAUGACUGUUGGAGUUGCCUCCCAAGAGAU
UCAGAAGCCAGACUUCAGGCAGCCUCAGGCUUUCUAGGUAUUGGUAUUGCCAGCUAGCUAAUADGUUCACAGCA
AAGGACUGAAGCUAGGUAUUAUGCAGAUUGGAAUAUAACUGGCCAGGUUCCUGGGAGUUUUGGAUACU
ACGACAUGAUGCCAGACCUUUGCUGACUGGGAGUAGAUUCUAAAUAUGAUGGUUACUGUGACAGUU
UGGAAAUUUGGCAGAUUGGUUAUAGCACAUUGCCUGCCUGAUAAGGACUGGCAGAGCAUUGUGUACUCCU
GUGAGUGCCUCUUUAUUGUGGCCUUUCAAAGCCAAUUAACAGAAAUCCGACAGUACUGCAAUCACUGGCG
AAUUUUGCUGACAUAUGAUAUCCUGGAAAGUAUAAGAUUUCUUGGACUGGACAUCUUUUAACCGAGAGAG
AAUUGUUGAUGUUGCUGGACCAGGGGUUGGAUUGACCAGAUAGUUAUGAUUGGCAACUUUGGCCUCAGCUG
GAUCAGCAAGUAACUCAGUUGGCCUCUGGGCUAUCUUGGCUUUAUUCUUGUUAUGUUAUGACCCCGACAC
AUCAGCCUUCAGCCAAAGCUCUCUUCAGGAUAAGGACGUAAUUGCCAUCAAGCACCCUUGGGCAAGCAAG
GGUACCAGCUUAGACAGGAGACAACUUUGAUGUGGGAACGACCUCUCAGGCUUAGCCUUGGGCUGUAGCUAU
GAUAAACCGGCAGGAGAUUGGUGGACCUUGCUUAUACCAUCCAGUUGCUUCCUGGGUAAGGAGUGGCCUGU
AAUCCUGCCUUCAUCAACAGCUCUCCUGUAAAGGAAGCUAGGGUUCUUGAUGGACUUCAGGUUA
GAAGUCACAUAUAUCCACAGGCACUGUUUGCUUCAGCUAGAAAUAACAUGCAGAUUGUCAUUAAGACUUAC
UUUAA (SEQ ID NO: 4)

FIG. 4

Human alpha-1 antitrypsin (A1AT) mRNA:

AUGCCGUCUUCUGUCUGGGGCAUCCUUGGAGCCUUGGAGCCUUGGUCUCCUUGUCUCCUUGGAGG
AUCCCAGGGAGUUGCUGCCAGACAGAUACAUCCACCAUGAUCAGGAUCACCCAACCUUCAACAAGAUAC
CCCCAACUUGGUGAGUUCGCCUUCAGCCUAUACCGCCAGCUGGCACACAGUCCACAGCACCACAUUUCUUCU
CCCCAGAGCAUCCGUACAGCCUUGCAUUGCUUCCUUGGGACCAAGGUCGACACUACAGGAUAUCCUGGA
GGCCUGAAUUUCAACCUACGGAGAUUCCGGAGGCUAGAUCCAUAGGCUUCCAGGACUCCUGGUACCCUC
AACCCAGCAGACGCCAGCUCAGCUGACACCGGCAUUGCCUUGUCCUAGCCAGGGCCUGAGCUAGUGGAUA
AGUUUUGGAGGAGUUAAAGUUGUACCAUCAGAGCCUUCACUGUCAUUCUGGGACACCGAAGAGGCCA
AGAAACAGAUACGAUUCGUGGAGAGGUACUCAAGGAAAUUGUGGAUUUGGUCAGGAGCUUGACAGAG
ACACAGUUUUGCUCUGGUAUUAUUAUUAAGGCAAUUGGAGAGACCCUUUGAAGUCAAGGACACCG
AGGAGAGGACUUCACGUGGACCGGACCCGUGAGGUGCCUAUGAUGAAGCGUUUAGGCAUGUUUAACA
UCCAGCACUGUAGAAGCUCAGCUGGGUGCUGAUGAAUACUUGGCAUCCACCGCCAUUCUUCUUCU
GCCUGAUGAGGGGAAACUACAGCACCGGAAAUGAACUCCACCAUUAUCAUCCAGUUCUGGAAAUGAA
GACAGAGGUCUGCCAGCUUACAUUUAACCAACUGUCCAUUACUGGAACCUAUGAUCUGAAGAGCGUCCUGGGUC
AACUGGCAUCACUAAGGUCUUCAGCAUUGGGGUGACCUUCUGGGGUCACAGAGGAGCCUUGAAGCUCUC
CAAGCCGUGCAUAAGGUCUGGUGACCAUUGCAGGAGAAAGGACUAGAGCUGGGGCAUGUUUUUAGAGGCC
AUACCCAUUCUAUCCCCCGAGGUCAGUUCACAAACCCUUGUCUUAUUGAUGAACAAAUACCAAGU
CUCCCCUUCUAUGGGAAAGUGGUGAUUCCACCCAAAUAA (SEQ ID NO: 5)

FIG. 5

Human Factor IX (FIX) mRNA:

AUGCAGCGGUGAACAUCAUGGAGGAGAUCAACAGGCGUCCUCCUUAUAGGAUAUCUCAGUG
CUGAAUGUACAGUUUUUUGAUCAGAAACGCCAACAAAUUCUGAGCGGAGAGGAGGUAUAUUCAGGUA
AAUUGGAAGAGUUUGUUCAGGGAACCUUGAGAGAGAUUGAAGAAAGUAGUUUUGAAGAGCACCAG
AAGUUUUUGAAACACUCGAAAGAACACUGAAUUUUGGAGGAGAUUGUAGGAGAUCCAGUGAGUCCCAUC
CAUGUUUAAAUUGGCGGAGUUUGGAGGAGCAUUAUUCCUAUGAUGUUGGUGUCCUUUGGAUUUGAAGGAA
AGAACUGUAUUAGAUUAACAUGUAACAUUAAGAAUGGCAGAUCCGAGCAGUUUUGUAATAAUAUGUCUGUA
ACAAGGUGUUUGCUCUGUACUGAGGGAUUGGACUUGCAGAAACCAGAGUCCUGUGAACCCAGCAGUGCCAUU
UCCAUGGGAAGAUUCUGUUUCACAAACUUUAAGCUCACCCUGUGAGGCUUUUUUCCUGAUGUGGACUUA
GUAUUUCUACUGAGCUGAAACCAUUUGGAUAACAUCACUCAAAGCACCACAAUUAUUAUGACUUCACUGGG
UUUGUGGAGAGAUCCCAACAGGUCUUUUUCCUUGGAGGUUUUGAUGGUAAGUUGAUGCAUUCU
GUGGAGCCUCUUCGUUAUAGAAAUAUGGAUUGUAACUGCUGCCACUGUUGAAACUGGUGUUAUUAUACAG
UUGUCGAGGUAACAUAUAUUGAGGAGACAGACAUAACAGAGCAAAGCAGAAUUGUAUUUAUUCUUC
ACCACAUCUACAUUGCAGCUUAUAUAGUACACCAUUGCCUUCUGGACUGGACGAAACCCUUAGUGCU
AAACAGCUACGUUACCUUUUGCAUUGCUGACAGGAUAACAGGAACAUCUUCCUCAAUUUGGAUCUGGCUAU
GUAAGUGCUGGGAGAGUCUCCACAAAGGGAGUCCAGUUUAGUCCUAGUACCUUAGAGUCCACUUGUU
GACCGAGCCACAUGUCUUGAUCUACAAAGUUCACCAUUAUAACAACAUUUUCUGGCUCCCAUGAAGGAG
GUAGAGUUCAUGUCAAGGAGAUAGUGGGGACCCCAUGUUAUGAAGUGGAGGGACCAUUAACUGGAA
UUUUAGCUGGGUGAGUGGCAUUAAGGCAAUUUGGAUAUAUACCAAGGUAUCCCGGUAUGUCAUCU
GGAUUAAGGAATAACAAGCUCUUA (SEQ ID NO: 6)

FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

HUMAN AGAL PROTEIN DETECTED IN MOUSE SERUM OVER TIME
AFTER TREATMENT VIA AGAL MRT




FIG. 12




FIG. 13




FIG. 14A




FIG. 14B




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22

HUMAN GLA PROTEIN DETECTED IN SERUM AFTER
TREATMENT VIA GLA MRT IN WT MICE USING HGT5001




FIG. 23A

HUMAN GAL PROTEIN PRODUCTION IN WT MOUSE TISSUE
24HR POST-ADMINISTRATION AFTER TREATMENT VIA GLA MRT USING
HGT5001




FIG. 23B




FIG. 24




FIG. 25




FIG. 26

LIPID NANOPARTICLE COMPOSITIONS AND METHODS FOR MRNA DELIVERY

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a divisional application of U.S. application Ser. No. 17/509,674 filed Oct. 25, 2021, issued as U.S. Pat. No. 11,291,734, which is a divisional application of U.S. application Ser. No. 17/341,016 filed Jun. 7, 2021, issued as U.S. Pat. No. 11,185,595, which is a divisional application of U.S. application Ser. No. 17/099,462 filed Nov. 16, 2020, issued as U.S. Pat. No. 11,052,159, which is divisional Application of U.S. application Ser. No. 16/931,231 filed Jul. 16, 2020, issued as U.S. Pat. No. 10,888,626, which is a divisional application of U.S. application Ser. No. 16/681,565 filed Nov. 12, 2019, now abandoned, which is a divisional application of U.S. application Ser. No. 16/502,672 filed Jul. 3, 2019, issued as U.S. Pat. No. 10,507,249, which is a divisional application of U.S. application Ser. No. 16/286,400 filed on Feb. 26, 2019, issued as U.S. Pat. No. 10,350,303 on Jul. 16, 2019, which is a divisional application of U.S. application Ser. No. 16/233,031 filed on Dec. 26, 2018, now issued as U.S. Pat. No. 10,413,618, which is a divisional application of U.S. application Ser. No. 15/482,117 filed on Apr. 7, 2017, now issued as U.S. Pat. No. 10,238,754, which is a divisional application of U.S. application Ser. No. 14/124,608 filed on Mar. 5, 2014, now abandoned, which is a U.S. National Entry claim priority to International Application PCT/US2012/041724 filed on Jun. 8, 2012, which claims priority to U.S. Provisional Application Ser. No. 61/494,881 filed Jun. 8, 2011, the disclosures of which are hereby incorporated by reference.

INCORPORATION-BY-REFERENCE OF SEQUENCE LISTING

This instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Dec. 17, 2021, is named MRT-1055US13_ST25.txt and is 10 KB in size. No new matter is hereby added.

Novel approaches and therapies are still needed for the treatment of protein and enzyme deficiencies. For example, lysosomal storage diseases are a group of approximately 50 rare inherited metabolic disorders that result from defects in lysosomal function, usually due to a deficiency of an enzyme required for metabolism. Fabry disease is a lysosomal storage disease that results from a deficiency of the enzyme alpha galactosidase (GLA), which causes a glycolipid known as globotriaosylceramide to accumulate in blood vessels and other tissues, leading to various painful manifestations. For certain diseases, like Fabry disease, there is a need for replacement of a protein or enzyme that is normally secreted by cells into the blood stream. Therapies, such as gene therapy, that increase the level or production of an affected protein or enzyme could provide a treatment or even a cure for such disorders. However, there have been several limitations to using conventional gene therapy for this purpose.

Conventional gene therapy involves the use of DNA for insertion of desired genetic information into host cells. The DNA introduced into the cell is usually integrated to a certain extent into the genome of one or more transfected cells, allowing for long-lasting action of the introduced

genetic material in the host. While there may be substantial benefits to such sustained action, integration of exogenous DNA into a host genome may also have many deleterious effects. For example, it is possible that the introduced DNA will be inserted into an intact gene, resulting in a mutation which impedes or even totally eliminates the function of the endogenous gene. Thus, gene therapy with DNA may result in the impairment of a vital genetic function in the treated host, such as e.g., elimination or deleteriously reduced production of an essential enzyme or interruption of a gene critical for the regulation of cell growth, resulting in unregulated or cancerous cell proliferation. In addition, with conventional DNA based gene therapy it is necessary for effective expression of the desired gene product to include a strong promoter sequence, which again may lead to undesirable changes in the regulation of normal gene expression in the cell. It is also possible that the DNA based genetic material will result in the induction of undesired anti-DNA antibodies, which in turn, may trigger a possibly fatal immune response. Gene therapy approaches using viral vectors can also result in an adverse immune response. In some circumstances, the viral vector may even integrate into the host genome. In addition, production of clinical grade viral vectors is also expensive and time consuming. Targeting delivery of the introduced genetic material using viral vectors can also be difficult to control. Thus, while DNA based gene therapy has been evaluated for delivery of secreted proteins using viral vectors (U.S. Pat. No. 6,066,626; US2004/0110709), these approaches may be limited for these various reasons.

Another obstacle apparent in these prior approaches at delivery of nucleic acids encoding secreted proteins, is in the levels of protein that are ultimately produced. It is difficult to achieve significant levels of the desired protein in the blood, and the amounts are not sustained over time. For example, the amount of protein produced by nucleic acid delivery does not reach normal physiological levels. See e.g., US2004/0110709.

In contrast to DNA, the use of RNA as a gene therapy agent is substantially safer because (1) RNA does not involve the risk of being stably integrated into the genome of the transfected cell, thus eliminating the concern that the introduced genetic material will disrupt the normal functioning of an essential gene, or cause a mutation that results in deleterious or oncogenic effects; (2) extraneous promoter sequences are not required for effective translation of the encoded protein, again avoiding possible deleterious side effects; (3) in contrast to plasmid DNA (pDNA), messenger RNA (mRNA) is devoid of immunogenic CpG motifs so that anti-RNA antibodies are not generated; and (4) any deleterious effects that do result from mRNA based on gene therapy would be of limited duration due to the relatively short half-life of RNA. In addition, it is not necessary for mRNA to enter the nucleus to perform its function, while DNA must overcome this major barrier.

One reason that mRNA based gene therapy has not been used more in the past is that mRNA is far less stable than DNA, especially when it reaches the cytoplasm of a cell and is exposed to degrading enzymes. The presence of a hydroxyl group on the second carbon of the sugar moiety in mRNA causes steric hinderance that prevents the mRNA from forming the more stable double helix structure of DNA and thus makes the mRNA more prone to hydrolytic degradation. As a result, until recently, it was widely believed that mRNA was too labile to withstand transfection protocols. Advances in RNA stabilizing modifications have sparked more interest in the use of mRNA in place of

plasmid DNA in gene therapy. Certain delivery vehicles, such as cationic lipid or polymer delivery vehicles may also help protect the transfected mRNA from endogenous RNases. Yet, in spite of increased stability of modified mRNA, delivery of mRNA to cells in vivo in a manner allowing for therapeutic levels of protein production is still a challenge, particularly for mRNA encoding full length proteins. While delivery of mRNA encoding secreted proteins has been contemplated (US2009/0286852), the levels of a full length secreted protein that would actually be produced via in vivo mRNA delivery are not known and there is not a reason to expect the levels would exceed those observed with DNA based gene therapy.

To date, significant progress using mRNA gene therapy has only been made in applications for which low levels of translation has not been a limiting factor, such as immunization with mRNA encoding antigens. Clinical trials involving vaccination against tumor antigens by intradermal injection of naked or protamine-complexed mRNA have demonstrated feasibility, lack of toxicity, and promising results. X. Su et al., *Mol. Pharmaceutics* 8:774-787 (2011). Unfortunately, low levels of translation has greatly restricted the exploitation of mRNA based gene therapy in other applications which require higher levels of sustained expression of the mRNA encoded protein to exert a biological or therapeutic effect.

The invention provides methods for delivery of mRNA gene therapeutic agents that lead to the production of therapeutically effective levels of secreted proteins via a "depot effect." In embodiments of the invention, mRNA encoding a secreted protein is loaded in lipid nanoparticles and delivered to target cells in vivo. Target cells then act as a depot source for production of soluble, secreted protein into the circulatory system at therapeutic levels. In some embodiments, the levels of secreted protein produced are above normal physiological levels.

The invention provides compositions and methods for intracellular delivery of mRNA in a liposomal transfer vehicle to one or more target cells for production of therapeutic levels of secreted functional protein.

The compositions and methods of the invention are useful in the management and treatment of a large number of diseases, in particular diseases which result from protein and/or enzyme deficiencies, wherein the protein or enzyme is normally secreted. Individuals suffering from such diseases may have underlying genetic defects that lead to the compromised expression of a protein or enzyme, including, for example, the non-synthesis of the secreted protein, the reduced synthesis of the secreted protein, or synthesis of a secreted protein lacking or having diminished biological activity. In particular, the methods and compositions of the invention are useful for the treatment of lysosomal storage disorders and/or the urea cycle metabolic disorders that occur as a result of one or more defects in the biosynthesis of secreted enzymes involved in the urea cycle.

The compositions of the invention comprise an mRNA, a transfer vehicle and, optionally, an agent to facilitate contact with, and subsequent transfection of a target cell. The mRNA can encode a clinically useful secreted protein. For example, the mRNA may encode a functional secreted urea cycle enzyme or a secreted enzyme implicated in lysosomal storage disorders. The mRNA can encode, for example, erythropoietin (e.g., human EPO) or α -galactosidase (e.g., human α -galactosidase (human GLA)).

In some embodiments the mRNA can comprise one or more modifications that confer stability to the mRNA (e.g., compared to a wild-type or native version of the mRNA) and

may also comprise one or more modifications relative to the wild-type which correct a defect implicated in the associated aberrant expression of the protein. For example, the nucleic acids of the present invention may comprise modifications to one or both of the 5' and 3' untranslated regions. Such modifications may include, but are not limited to, the inclusion of a partial sequence of a cytomegalovirus (CMV) immediate-early 1 (IE1) gene, a poly A tail, a Cap1 structure or a sequence encoding human growth hormone (hGH)). In some embodiments, the mRNA is modified to decrease mRNA immunogenicity.

Methods of treating a subject comprising administering a composition of the invention, are also contemplated. For example, methods of treating or preventing conditions in which production of a particular secreted protein and/or utilization of a particular secreted protein is inadequate or compromised are provided. In one embodiment, the methods provided herein can be used to treat a subject having a deficiency in one or more urea cycle enzymes or in one or more enzymes deficient in a lysosomal storage disorder.

In a preferred embodiment, the mRNA in the compositions of the invention is formulated in a liposomal transfer vehicle to facilitate delivery to the target cell. Contemplated transfer vehicles may comprise one or more cationic lipids, non-cationic lipids, and/or PEG-modified lipids. For example, the transfer vehicle may comprise at least one of the following cationic lipids: C12-200, DLin-KC2-DMA, DODAP, HGT4003, ICE, HGT5000, or HGT5001. In embodiments, the transfer vehicle comprises cholesterol (chol) and/or a PEG-modified lipid. In some embodiments, the transfer vehicles comprises DMG-PEG2K. In certain embodiments, the transfer vehicle comprises one of the following lipid formulations: C12-200, DOPE, chol, DMG-PEG2K; DODAP, DOPE, cholesterol, DMG-PEG2K; HGT5000, DOPE, chol, DMG-PEG2K, HGT5001, DOPE, chol, DMG-PEG2K.

The invention also provides compositions and methods useful for facilitating the transfection and delivery of one or more mRNA molecules to target cells capable of exhibiting the "depot effect." For example, the compositions and methods of the present invention contemplate the use of targeting ligands capable of enhancing the affinity of the composition to one or more target cells. In one embodiment, the targeting ligand is apolipoprotein-B or apolipoprotein-E and corresponding target cells express low-density lipoprotein receptors, thereby facilitating recognition of the targeting ligand. The methods and compositions of the present invention may be used to preferentially target a vast number of target cells. For example, contemplated target cells include, but are not limited to, hepatocytes, epithelial cells, hematopoietic cells, epithelial cells, endothelial cells, lung cells, bone cells, stem cells, mesenchymal cells, neural cells, cardiac cells, adipocytes, vascular smooth muscle cells, cardiomyocytes, skeletal muscle cells, beta cells, pituitary cells, synovial lining cells, ovarian cells, testicular cells, fibroblasts, B cells, T cells, reticulocytes, leukocytes, granulocytes and tumor cells.

In embodiments, the secreted protein is produced by the target cell for sustained amounts of time. For example, the secreted protein may be produced for more than one hour, more than four, more than six, more than 12, more than 24, more than 48 hours, or more than 72 hours after administration. In some embodiments the polypeptide is expressed at a peak level about six hours after administration. In some embodiments the expression of the polypeptide is sustained at least at a therapeutic level. In some embodiments the polypeptide is expressed at at least a therapeutic level for

more than one, more than four, more than six, more than 12, more than 24, more than 48 hours, or more than 72 hours after administration. In some embodiments the polypeptide is detectable at the level in patient serum or tissue (e.g., liver, or lung). In some embodiments, the level of detectable polypeptide is from continuous expression from the mRNA composition over periods of time of more than one, more than four, more than six, more than 12, more than 24, more than 48 hours, or more than 72 hours after administration.

In certain embodiments, the secreted protein is produced at levels above normal physiological levels. The level of secreted protein may be increased as compared to a control.

In some embodiments the control is the baseline physiological level of the polypeptide in a normal individual or in a population of normal individuals. In other embodiments the control is the baseline physiological level of the polypeptide in an individual having a deficiency in the relevant protein or polypeptide or in a population of individuals having a deficiency in the relevant protein or polypeptide. In some embodiments the control can be the normal level of the relevant protein or polypeptide in the individual to whom the composition is administered. In other embodiments the control is the expression level of the polypeptide upon other therapeutic intervention, e.g., upon direct injection of the corresponding polypeptide, at one or more comparable time points.

In certain embodiments the polypeptide is expressed by the target cell at a level which is at least 1.5-fold, at least 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, 30-fold, at least 100-fold, at least 500-fold, at least 5000-fold, at least 50,000-fold or at least 100,000-fold greater than a control. In some embodiments, the fold increase of expression greater than control is sustained for more than one, more than four, more than six, more than 12, more than 24, or more than 48 hours, or more than 72 hours after administration. For example, in one embodiment, the levels of secreted protein are detected in the serum at least 1.5-fold, at least 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, 30-fold, at least 100-fold, at least 500-fold, at least 5000-fold, at least 50,000-fold or at least 100,000-fold greater than a control for at least 48 hours or 2 days. In certain embodiments, the levels of secreted protein are detectable at 3 days, 4 days, 5 days, or 1 week or more after administration. Increased levels of secreted protein may be observed in the serum and/or in a tissue (e.g. liver, lung).

In some embodiments, the method yields a sustained circulation half-life of the desired secreted protein. For example, the secreted protein may be detected for hours or days longer than the half-life observed via subcutaneous injection of the secreted protein. In embodiments, the half-life of the secreted protein is sustained for more than 1 day, 2 days, 3 days, 4 days, 5 days, or 1 week or more.

In some embodiments administration comprises a single or repeated doses. In certain embodiments, the dose is administered intravenously, or by pulmonary delivery.

The polypeptide can be, for example, one or more of erythropoietin, α -galactosidase, LDL receptor, Factor VIII, Factor IX, α -L-iduronidase (for MPS I), iduronate sulfatase (for MPS II), heparin-N-sulfatase (for MPS IIIA), α -N-acetylglucosaminidase (for MPS IIIB), galactose 6-sulfatase (for MPS IVA), lysosomal acid lipase, arylsulfatase-A.

Certain embodiments relate to compositions and methods that provide to a cell or subject mRNA, at least a part of which encodes a functional protein, in an amount that is substantially less than the amount of corresponding functional protein generated from that mRNA. Put another way, in certain embodiments the mRNA delivered to the cell can

produce an amount of protein that is substantially greater than the amount of mRNA delivered to the cell. For example, in a given amount of time, for example 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 20, or 24 hours from administration of the mRNA to a cell or subject, the amount of corresponding protein generated by that mRNA can be at least 1.5, 2, 3, 5, 10, 15, 20, 25, 50, 100, 150, 200, 250, 300, 400, 500, or more times greater than the amount of mRNA actually administered to the cell or subject. This can be measured on a mass-by-mass basis, on a mole-by-mole basis, and/or on a molecule-by-molecule basis. The protein can be measured in various ways. For example, for a cell, the measured protein can be measured as intracellular protein, extracellular protein, or a combination of the two. For a subject, the measured protein can be protein measured in serum; in a specific tissue or tissues such as the liver, kidney, heart, or brain; in a specific cell type such as one of the various cell types of the liver or brain; or in any combination of serum, tissue, and/or cell type. Moreover, a baseline amount of endogenous protein can be measured in the cell or subject prior to administration of the mRNA and then subtracted from the protein measured after administration of the mRNA to yield the amount of corresponding protein generated from the mRNA. In this way, the mRNA can provide a reservoir or depot source of a large amount of therapeutic material to the cell or subject, for example, as compared to amount of mRNA delivered to the cell or subject. The depot source can act as a continuous source for polypeptide expression from the mRNA over sustained periods of time.

The above discussed and many other features and attendant advantages of the present invention will become better understood by reference to the following detailed description of the invention when taken in conjunction with the accompanying examples. The various embodiments described herein are complimentary and can be combined or used together in a manner understood by the skilled person in view of the teachings contained herein.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the nucleotide sequence of a 5' CMV sequence (SEQ ID NO:1), wherein X, if present is GGA.

FIG. 2 shows the nucleotide sequence of a 3' hGH sequence (SEQ ID NO:2).

FIG. 3 shows the nucleotide sequence of human erythropoietic (EPO) mRNA (SEQ ID NO:3). This sequence can be flanked on the 5' end with SEQ ID NO:1 and on the 3' end with SEQ ID NO:2.

FIG. 4 shows the nucleotide sequence of human alpha-galactosidase (GLA) mRNA (SEQ ID NO:4). This sequence can be flanked on the 5' end with SEQ ID NO:1 and on the 3' end with SEQ ID NO:2.

FIG. 5 shows the nucleotide sequence of human alpha-1 antitrypsin (A1AT) mRNA (SEQ ID NO:5). This sequence can be flanked on the 5' end with SEQ ID NO:1 and on the 3' end with SEQ ID NO:2.

FIG. 6 shows the nucleotide sequence of human factor IX (FIX) mRNA (SEQ ID NO:6). This sequence can be flanked on the 5' end with SEQ ID NO:1 and on the 3' end with SEQ ID NO:2.

FIG. 7 shows quantification of secreted hEPO protein levels as measured via ELISA. The protein detected is a result of its production from hEPO mRNA delivered intravenously via a single dose of various lipid nanoparticle formulations. The formulations C12-200 (30 ug), HGT4003 (150 ug), ICE (100 ug), DODAP (200 ug) are represented as

the cationic/ionizable lipid component of each test article (Formulations 1-4). Values are based on blood sample four hours post-administration.

FIG. 8 shows the hematocrit measurement of mice treated with a single IV dose of human EPO mRNA-loaded lipid nanoparticles (Formulations 1-4). Whole blood samples were taken at 4 hr (Day 1), 24 hr (Day 2), 4 days, 7 days, and 10 days post-administration.

FIG. 9 shows hematocrit measurements of mice treated with human EPO-mRNA-loaded lipid nanoparticles with either a single IV dose or three injections (day 1, day 3, day 5). Whole blood samples were taken prior to injection (day -4), day 7, and day 15. Formulation 1 was administered: (30 ug, single dose) or (3×10 ug, dose day 1, day 3, day 5); Formulation 2 was administered: (3×50 ug, dose day 1, day 3, day 5).

FIG. 10 shows quantification of secreted human α -galactosidase (hGLA) protein levels as measured via ELISA. The protein detected is a result of the production from hGLA mRNA delivered via lipid nanoparticles (Formulation 1; 30 ug single intravenous dose, based on encapsulated mRNA). hGLA protein is detected through 48 hours.

FIG. 11 shows hGLA activity in serum. hGLA activity was measured using substrate 4-methylumbelliferyl- α -D-galactopyranoside (4-MU- α -gal) at 37° C. Data are average of 6 to 9 individual measurements.

FIG. 12 shows quantification of hGLA protein levels in serum as measured via ELISA. Protein is produced from hGLA mRNA delivered via C12-200-based lipid nanoparticles (C12-200:DOPE:Chol:DMGPEG2K, 40:30:25:5 (Formulation 1); 30 ug mRNA based on encapsulated mRNA, single IV dose). hGLA protein is monitored through 72 hours. per single intravenous dose, based on encapsulated mRNA). hGLA protein is monitored through 72 hours.

FIG. 13 shows quantification of hGLA protein levels in liver, kidney, and spleen as measured via ELISA. Protein is produced from hGLA mRNA delivered via C12-200-based lipid nanoparticles (Formulation 1; 30 ug mRNA based on encapsulated mRNA, single IV dose). hGLA protein is monitored through 72 hours.

FIG. 14A shows a dose response study monitoring protein production of hGLA as secreted MRT-derived human GLA protein in serum. Samples were measured 24 hours post-administration (Formulation 1; single dose, IV, N=4 mice/group) and quantified via ELISA.

FIG. 14B shows a dose response study monitoring protein production of hGLA as secreted MRT-derived human GLA protein in liver. Samples were measured 24 hours post-administration (Formulation 1; single dose, IV, N=4 mice/group) and quantified via ELISA.

FIG. 15 shows the pharmacokinetic profiles of ERT-based Alpha-galactosidase in athymic nude mice (40 ug/kg dose) and hGLA protein produced from MRT (Formulation 1; 1.0 mg/kg mRNA dose).

FIG. 16 shows the quantification of secreted hGLA protein levels in MRT-treated Fabry mice as measured using ELISA. hGLA protein is produced from hGLA mRNA delivered via C12-200-based lipid nanoparticles (Formulation 1; 10 ug mRNA per single intravenous dose, based on encapsulated mRNA). Serum is monitored through 72 hours.

FIG. 17 shows the quantification of hGLA protein levels in liver, kidney, spleen, and heart of MRT-treated Fabry KO mice as measured via ELISA. Protein is produced from hGLA mRNA delivered via C12-200-based lipid nanoparticles (Formulation 1; 30 ug mRNA based on encapsulated mRNA, single IV dose). hGLA protein is monitored through

72 hours. Literature values representing normal physiological levels are graphed as dashed lines.

FIG. 18 shows the quantification of secreted hGLA protein levels in MRT and Alpha-galactosidase-treated Fabry mice as measured using ELISA. Both therapies were dosed as a single 1.0 mg/kg intravenous dose.

FIG. 19 shows the quantification of hGLA protein levels in liver, kidney, spleen, and heart of MRT and ERT (Alpha-galactosidase)-treated Fabry KO mice as measured via ELISA. Protein produced from hGLA mRNA delivered via lipid nanoparticles (Formulation 1; 1.0 mg/kg mRNA based on encapsulated mRNA, single IV dose).

FIG. 20 shows the relative quantification of globotri- asylceramide (Gb3) and lyso-Gb3 in the kidneys of treated and untreated mice. Male Fabry KO mice were treated with a single dose either GLA mRNA-loaded lipid nanoparticles or Alpha-galactosidase at 1.0 mg/kg. Amounts reflect quantity of Gb3/lyso-Gb3 one week post-administration.

FIG. 21 shows the relative quantification of globotri- asylceramide (Gb3) and lyso-Gb3 in the heart of treated and untreated mice. Male Fabry KO mice were treated with a single dose either GLA mRNA-loaded lipid nanoparticles or Alpha-galactosidase at 1.0 mg/kg. Amounts reflect quantity of Gb3/lyso-Gb3 one week post-administration.

FIG. 22 shows a dose response study monitoring protein production of GLA as secreted MRT-derived human GLA protein in serum. Samples were measured 24 hours post-administration (single dose, IV, N=4 mice/group) of either HGT4003 (Formulation 3) or HGT5000-based lipid nanoparticles (Formulation 5) and quantified via ELISA.

FIG. 23A shows hGLA protein production as measured in serum. Samples were measured 6 hours and 24 hours post-administration (single dose, IV, N=4 mice/group) of HGT5001-based lipid nanoparticles (Formulation 6) and quantified via ELISA.

FIG. 23B shows hGLA protein production as measured in liver, kidney, and spleen (B). Samples were measured 6 hours and 24 hours post-administration (single dose, IV, N=4 mice/group) of HGT5001-based lipid nanoparticles (Formulation 6) and quantified via ELISA.

FIG. 24 shows the quantification of secreted human Factor IX protein levels measured using ELISA (mean ng/mL \pm standard deviation). FIX protein is produced from FIX mRNA delivered via C12-200-based lipid nanoparticles (C12-200:DOPE:Chol:DMGPEG2K, 40:30:25:5 (Formulation 1); 30 ug mRNA per single intravenous dose, based on encapsulated mRNA). FIX protein is monitored through 72 hours. (n=24 mice)

FIG. 25 shows the quantification of secreted human α -1-antitrypsin (A1AT) protein levels measured using ELISA. A1AT protein is produced from A1AT mRNA delivered via C12-200-based lipid nanoparticles (C12-200:DOPE:Chol:DMGPEG2K, 40:30:25:5 (Formulation 1); 30 ug mRNA per single intravenous dose, based on encapsulated mRNA). A1AT protein is monitored through 24 hours.

FIG. 26 shows an ELISA-based quantification of hEPO protein detected in the lungs and serum of treated mice after intratracheal administration of hEPO mRNA-loaded nanoparticles (measured mIU) (C12-200, HGT5000, or HGT5001-based lipid nanoparticles; Formulations 1, 5, 6 respectively). Animals were sacrificed 6 hours post-administration (n=4 mice per group).

DESCRIPTION OF EXEMPLARY EMBODIMENTS

The invention provides compositions and methods for intracellular delivery of mRNA in a liposomal transfer

vehicle to one or more target cells for production of therapeutic levels of secreted functional protein.

The term “functional,” as used herein to qualify a protein or enzyme, means that the protein or enzyme has biological activity, or alternatively is able to perform the same, or a similar function as the native or normally-functioning protein or enzyme. The mRNA compositions of the invention are useful for the treatment of a various metabolic or genetic disorders, and in particular those genetic or metabolic disorders which involve the non-expression, misexpression or deficiency of a protein or enzyme. The term “therapeutic levels” refers to levels of protein detected in the blood or tissues that are above control levels, wherein the control may be normal physiological levels, or the levels in the subject prior to administration of the mRNA composition. The term “secreted” refers to protein that is detected outside the target cell, in extracellular space. The protein may be detected in the blood or in tissues. In the context of the present invention the term “produced” is used in its broadest sense to refer the translation of at least one mRNA into a protein or enzyme. As provided herein, the compositions include a transfer vehicle. As used herein, the term “transfer vehicle” includes any of the standard pharmaceutical carriers, diluents, excipients and the like which are generally intended for use in connection with the administration of biologically active agents, including nucleic acids. The compositions and in particular the transfer vehicles described herein are capable of delivering mRNA to the target cell. In embodiments, the transfer vehicle is a lipid nanoparticle.

mRNA

The mRNA in the compositions of the invention may encode, for example, a secreted hormone, enzyme, receptor, polypeptide, peptide or other protein of interest that is normally secreted. In one embodiment of the invention, the mRNA may optionally have chemical or biological modifications which, for example, improve the stability and/or half-life of such mRNA or which improve or otherwise facilitate protein production.

The methods of the invention provide for optional co-delivery of one or more unique mRNA to target cells, for example, by combining two unique mRNAs into a single transfer vehicle. In one embodiment of the present invention, a therapeutic first mRNA, and a therapeutic second mRNA, may be formulated in a single transfer vehicle and administered. The present invention also contemplates co-delivery and/or co-administration of a therapeutic first mRNA and a second nucleic acid to facilitate and/or enhance the function or delivery of the therapeutic first mRNA. For example, such a second nucleic acid (e.g., exogenous or synthetic mRNA) may encode a membrane transporter protein that upon expression (e.g., translation of the exogenous or synthetic mRNA) facilitates the delivery or enhances the biological activity of the first mRNA. Alternatively, the therapeutic first mRNA may be administered with a second nucleic acid that functions as a “chaperone” for example, to direct the folding of either the therapeutic first mRNA.

The methods of the invention also provide for the delivery of one or more therapeutic nucleic acids to treat a single disorder or deficiency, wherein each such therapeutic nucleic acid functions by a different mechanism of action. For example, the compositions of the present invention may comprise a therapeutic first mRNA which, for example, is administered to correct an endogenous protein or enzyme deficiency, and which is accompanied by a second nucleic acid, which is administered to deactivate or “knock-down” a malfunctioning endogenous nucleic acid and its protein or

enzyme product. Such “second” nucleic acids may encode, for example mRNA or siRNA.

Upon transfection, a natural mRNA in the compositions of the invention may decay with a half-life of between 30 minutes and several days. The mRNA in the compositions of the invention preferably retain at least some ability to be translated, thereby producing a functional secreted protein or enzyme. Accordingly, the invention provides compositions comprising and methods of administering a stabilized mRNA. In some embodiments of the invention, the activity of the mRNA is prolonged over an extended period of time. For example, the activity of the mRNA may be prolonged such that the compositions of the present invention are administered to a subject on a semi-weekly or bi-weekly basis, or more preferably on a monthly, bi-monthly, quarterly or an annual basis. The extended or prolonged activity of the mRNA of the present invention, is directly related to the quantity of secreted functional protein or enzyme produced from such mRNA. Similarly, the activity of the compositions of the present invention may be further extended or prolonged by modifications made to improve or enhance translation of the mRNA. Furthermore, the quantity of functional protein or enzyme produced by the target cell is a function of the quantity of mRNA delivered to the target cells and the stability of such mRNA. To the extent that the stability of the mRNA of the present invention may be improved or enhanced, the half-life, the activity of the produced secreted protein or enzyme and the dosing frequency of the composition may be further extended.

Accordingly, in some embodiments, the mRNA in the compositions of the invention comprise at least one modification which confers increased or enhanced stability to the nucleic acid, including, for example, improved resistance to nuclease digestion *in vivo*. As used herein, the terms “modification” and “modified” as such terms relate to the nucleic acids provided herein, include at least one alteration which preferably enhances stability and renders the mRNA more stable (e.g., resistant to nuclease digestion) than the wild-type or naturally occurring version of the mRNA. As used herein, the terms “stable” and “stability” as such terms relate to the nucleic acids of the present invention, and particularly with respect to the mRNA, refer to increased or enhanced resistance to degradation by, for example nucleases (i.e., endonucleases or exonucleases) which are normally capable of degrading such mRNA. Increased stability can include, for example, less sensitivity to hydrolysis or other destruction by endogenous enzymes (e.g., endonucleases or exonucleases) or conditions within the target cell or tissue, thereby increasing or enhancing the residence of such mRNA in the target cell, tissue, subject and/or cytoplasm. The stabilized mRNA molecules provided herein demonstrate longer half-lives relative to their naturally occurring, unmodified counterparts (e.g. the wild-type version of the mRNA). Also contemplated by the terms “modification” and “modified” as such terms related to the mRNA of the present invention are alterations which improve or enhance translation of mRNA nucleic acids, including for example, the inclusion of sequences which function in the initiation of protein translation (e.g., the Kozac consensus sequence). (Kozak, M., *Nucleic Acids Res* 15 (20): 8125-48 (1987)).

In some embodiments, the mRNA of the invention have undergone a chemical or biological modification to render them more stable. Exemplary modifications to an mRNA include the depletion of a base (e.g., by deletion or by the substitution of one nucleotide for another) or modification of a base, for example, the chemical modification of a base. The phrase “chemical modifications” as used herein, includes

modifications which introduce chemistries which differ from those seen in naturally occurring mRNA, for example, covalent modifications such as the introduction of modified nucleotides, (e.g., nucleotide analogs, or the inclusion of pendant groups which are not naturally found in such mRNA molecules).

In addition, suitable modifications include alterations in one or more nucleotides of a codon such that the codon encodes the same amino acid but is more stable than the codon found in the wild-type version of the mRNA. For example, an inverse relationship between the stability of RNA and a higher number cytidines (C's) and/or uridines (U's) residues has been demonstrated, and RNA devoid of C and U residues have been found to be stable to most RNases (Heidenreich, et al. *J Biol Chem* 269, 2131-8 (1994)). In some embodiments, the number of C and/or U residues in an mRNA sequence is reduced. In a another embodiment, the number of C and/or U residues is reduced by substitution of one codon encoding a particular amino acid for another codon encoding the same or a related amino acid. Contemplated modifications to the mRNA nucleic acids of the present invention also include the incorporation of pseudouridines. The incorporation of pseudouridines into the mRNA nucleic acids of the present invention may enhance stability and translational capacity, as well as diminishing immunogenicity in vivo. See, e.g., Karikó, K., et al., *Molecular Therapy* 16 (11): 1833-1840 (2008). Substitutions and modifications to the mRNA of the present invention may be performed by methods readily known to one of ordinary skill in the art.

The constraints on reducing the number of C and U residues in a sequence will likely be greater within the coding region of an mRNA, compared to an untranslated region, (i.e., it will likely not be possible to eliminate all of the C and U residues present in the message while still retaining the ability of the message to encode the desired amino acid sequence). The degeneracy of the genetic code, however presents an opportunity to allow the number of C and/or U residues that are present in the sequence to be reduced, while maintaining the same coding capacity (i.e., depending on which amino acid is encoded by a codon, several different possibilities for modification of RNA sequences may be possible). For example, the codons for Gly can be altered to GGA or GGG instead of GGU or GGC.

The term modification also includes, for example, the incorporation of non-nucleotide linkages or modified nucleotides into the mRNA sequences of the present invention (e.g., modifications to one or both the 3' and 5' ends of an mRNA molecule encoding a functional secreted protein or enzyme). Such modifications include the addition of bases to an mRNA sequence (e.g., the inclusion of a poly A tail or a longer poly A tail), the alteration of the 3' UTR or the 5' UTR, complexing the mRNA with an agent (e.g., a protein or a complementary nucleic acid molecule), and inclusion of elements which change the structure of an mRNA molecule (e.g., which form secondary structures).

The poly A tail is thought to stabilize natural messengers. Therefore, in one embodiment a long poly A tail can be added to an mRNA molecule thus rendering the mRNA more stable. Poly A tails can be added using a variety of art-recognized techniques. For example, long poly A tails can be added to synthetic or in vitro transcribed mRNA using poly A polymerase (Yokoe, et al. *Nature Biotechnology*. 1996; 14: 1252-1256). A transcription vector can also encode long poly A tails. In addition, poly A tails can be added by transcription directly from PCR products. In one embodiment, the length of the poly A tail is at least about 90,

200, 300, 400 at least 500 nucleotides. In one embodiment, the length of the poly A tail is adjusted to control the stability of a modified mRNA molecule of the invention and, thus, the transcription of protein. For example, since the length of the poly A tail can influence the half-life of an mRNA molecule, the length of the poly A tail can be adjusted to modify the level of resistance of the mRNA to nucleases and thereby control the time course of protein expression in a cell. In one embodiment, the stabilized mRNA molecules are sufficiently resistant to in vivo degradation (e.g., by nucleases), such that they may be delivered to the target cell without a transfer vehicle.

In one embodiment, an mRNA can be modified by the incorporation 3' and/or 5' untranslated (UTR) sequences which are not naturally found in the wild-type mRNA. In one embodiment, 3' and/or 5' flanking sequence which naturally flanks an mRNA and encodes a second, unrelated protein can be incorporated into the nucleotide sequence of an mRNA molecule encoding a therapeutic or functional protein in order to modify it. For example, 3' or 5' sequences from mRNA molecules which are stable (e.g., globin, actin, GAPDH, tubulin, histone, or citric acid cycle enzymes) can be incorporated into the 3' and/or 5' region of a sense mRNA nucleic acid molecule to increase the stability of the sense mRNA molecule. See, e.g., US2003/0083272.

In some embodiments, the mRNA in the compositions of the invention include modification of the 5' end of the mRNA to include a partial sequence of a CMV immediate-early 1 (IE1) gene, or a fragment thereof (e.g., SEQ ID NO:1) to improve the nuclease resistance and/or improve the half-life of the mRNA. In addition to increasing the stability of the mRNA nucleic acid sequence, it has been surprisingly discovered the inclusion of a partial sequence of a CMV immediate-early 1 (IE1) gene enhances the translation of the mRNA and the expression of the functional protein or enzyme. Also contemplated is the inclusion of a human growth hormone (hGH) gene sequence, or a fragment thereof (e.g., SEQ ID NO:2) to the 3' ends of the nucleic acid (e.g., mRNA) to further stabilize the mRNA. Generally, preferred modifications improve the stability and/or pharmacokinetic properties (e.g., half-life) of the mRNA relative to their unmodified counterparts, and include, for example modifications made to improve such mRNA's resistance to in vivo nuclease digestion.

Further contemplated are variants of the nucleic acid sequence of SEQ ID NO:1 and/or SEQ ID NO:2, wherein the variants maintain the functional properties of the nucleic acids including stabilization of the mRNA and/or pharmacokinetic properties (e.g., half-life). Variants may have greater than 90%, greater than 95%, greater than 98%, or greater than 99% sequence identity to SEQ ID NO:1 or SEQ ID NO:2.

In some embodiments, the composition can comprise a stabilizing reagent. The compositions can include one or more formulation reagents that bind directly or indirectly to, and stabilize the mRNA, thereby enhancing residence time in the target cell. Such reagents preferably lead to an improved half-life of the mRNA in the target cells. For example, the stability of an mRNA and efficiency of translation may be increased by the incorporation of "stabilizing reagents" that form complexes with the mRNA that naturally occur within a cell (see e.g., U.S. Pat. No. 5,677,124). Incorporation of a stabilizing reagent can be accomplished for example, by combining the poly A and a protein with the mRNA to be stabilized in vitro before loading or encapsulating the mRNA within a transfer vehicle. Exemplary stabilizing reagents include one or more proteins, peptides,

aptamers, translational accessory protein, mRNA binding proteins, and/or translation initiation factors.

Stabilization of the compositions may also be improved by the use of opsonization-inhibiting moieties, which are typically large hydrophilic polymers that are chemically or physically bound to the transfer vehicle (e.g., by the intercalation of a lipid-soluble anchor into the membrane itself, or by binding directly to active groups of membrane lipids). These opsonization-inhibiting hydrophilic polymers form a protective surface layer which significantly decreases the uptake of the liposomes by the macrophage-monocyte system and reticulo-endothelial system (e.g., as described in U.S. Pat. No. 4,920,016, the entire disclosure of which is herein incorporated by reference). Transfer vehicles modified with opsonization-inhibition moieties thus remain in the circulation much longer than their unmodified counterparts.

When RNA is hybridized to a complementary nucleic acid molecule (e.g., DNA or RNA) it may be protected from nucleases. (Krieg, et al. *Methods in Enzymology*. 1987; 155, 397-415). The stability of hybridized mRNA is likely due to the inherent single strand specificity of most RNases. In some embodiments, the stabilizing reagent selected to complex a mRNA is a eukaryotic protein, (e.g., a mammalian protein). In yet another embodiment, the mRNA can be modified by hybridization to a second nucleic acid molecule. If an entire mRNA molecule were hybridized to a complementary nucleic acid molecule translation initiation may be reduced. In some embodiments the 5' untranslated region and the AUG start region of the mRNA molecule may optionally be left unhybridized. Following translation initiation, the unwinding activity of the ribosome complex can function even on high affinity duplexes so that translation can proceed. (Liebhaber. *J. Mol. Biol.* 1992; 226: 2-13; Monia, et al. *J Biol Chem.* 1993; 268: 14514-22.)

It will be understood that any of the above described methods for enhancing the stability of mRNA may be used either alone or in combination with one or more of any of the other above-described methods and/or compositions.

The mRNA of the present invention may be optionally combined with a reporter gene (e.g., upstream or downstream of the coding region of the mRNA) which, for example, facilitates the determination of mRNA delivery to the target cells or tissues. Suitable reporter genes may include, for example, Green Fluorescent Protein mRNA (GFP mRNA), *Renilla* Luciferase mRNA (Luciferase mRNA), Firefly Luciferase mRNA, or any combinations thereof. For example, GFP mRNA may be fused with a mRNA encoding a secretable protein to facilitate confirmation of mRNA localization in the target cells that will act as a depot for protein production.

As used herein, the terms "transfect" or "transfection" mean the intracellular introduction of a mRNA into a cell, or preferably into a target cell. The introduced mRNA may be stably or transiently maintained in the target cell. The term "transfection efficiency" refers to the relative amount of mRNA taken up by the target cell which is subject to transfection. In practice, transfection efficiency is estimated by the amount of a reporter nucleic acid product expressed by the target cells following transfection. Preferred embodiments include compositions with high transfection efficacies and in particular those compositions that minimize adverse effects which are mediated by transfection of non-target cells. The compositions of the present invention that demonstrate high transfection efficacies improve the likelihood that appropriate dosages of the mRNA will be delivered to the target cell, while minimizing potential systemic adverse effects. In one embodiment of the present invention, the

transfer vehicles of the present invention are capable of delivering large mRNA sequences (e.g., mRNA of at least 1 kDa, 1.5 kDa, 2 kDa, 2.5 kDa, 5 kDa, 10 kDa, 12 kDa, 15 kDa, 20 kDa, 25 kDa, 30 kDa, or more). The mRNA can be formulated with one or more acceptable reagents, which provide a vehicle for delivering such mRNA to target cells. Appropriate reagents are generally selected with regard to a number of factors, which include, among other things, the biological or chemical properties of the mRNA, the intended route of administration, the anticipated biological environment to which such mRNA will be exposed and the specific properties of the intended target cells. In some embodiments, transfer vehicles, such as liposomes, encapsulate the mRNA without compromising biological activity. In some embodiments, the transfer vehicle demonstrates preferential and/or substantial binding to a target cell relative to non-target cells. In a preferred embodiment, the transfer vehicle delivers its contents to the target cell such that the mRNA are delivered to the appropriate subcellular compartment, such as the cytoplasm.

Transfer Vehicle

In embodiments, the transfer vehicle in the compositions of the invention is a liposomal transfer vehicle, e.g. a lipid nanoparticle. In one embodiment, the transfer vehicle may be selected and/or prepared to optimize delivery of the mRNA to a target cell. For example, if the target cell is a hepatocyte the properties of the transfer vehicle (e.g., size, charge and/or pH) may be optimized to effectively deliver such transfer vehicle to the target cell, reduce immune clearance and/or promote retention in that target cell. Alternatively, if the target cell is the central nervous system (e.g., mRNA administered for the treatment of neurodegenerative diseases may specifically target brain or spinal tissue), selection and preparation of the transfer vehicle must consider penetration of, and retention within the blood brain barrier and/or the use of alternate means of directly delivering such transfer vehicle to such target cell. In one embodiment, the compositions of the present invention may be combined with agents that facilitate the transfer of exogenous mRNA (e.g., agents which disrupt or improve the permeability of the blood brain barrier and thereby enhance the transfer of exogenous mRNA to the target cells).

The use of liposomal transfer vehicles to facilitate the delivery of nucleic acids to target cells is contemplated by the present invention. Liposomes (e.g., liposomal lipid nanoparticles) are generally useful in a variety of applications in research, industry, and medicine, particularly for their use as transfer vehicles of diagnostic or therapeutic compounds in vivo (Lasic, *Trends Biotechnol.*, 16: 307-321, 1998; Drummond et al., *Pharmacol. Rev.*, 51: 691-743, 1999) and are usually characterized as microscopic vesicles having an interior aqueous space sequestered from an outer medium by a membrane of one or more bilayers. Bilayer membranes of liposomes are typically formed by amphiphilic molecules, such as lipids of synthetic or natural origin that comprise spatially separated hydrophilic and hydrophobic domains (Lasic, *Trends Biotechnol.*, 16: 307-321, 1998). Bilayer membranes of the liposomes can also be formed by amphiphilic polymers and surfactants (e.g., polymerosomes, niosomes, etc.).

In the context of the present invention, a liposomal transfer vehicle typically serves to transport the mRNA to the target cell. For the purposes of the present invention, the liposomal transfer vehicles are prepared to contain the desired nucleic acids. The process of incorporation of a desired entity (e.g., a nucleic acid) into a liposome is often referred to as "loading" (Lasic, et al., *FEBS Lett.*, 312:

255-258, 1992). The liposome-incorporated nucleic acids may be completely or partially located in the interior space of the liposome, within the bilayer membrane of the liposome, or associated with the exterior surface of the liposome membrane. The incorporation of a nucleic acid into liposomes is also referred to herein as “encapsulation” wherein the nucleic acid is entirely contained within the interior space of the liposome. The purpose of incorporating a mRNA into a transfer vehicle, such as a liposome, is often to protect the nucleic acid from an environment which may contain enzymes or chemicals that degrade nucleic acids and/or systems or receptors that cause the rapid excretion of the nucleic acids. Accordingly, in a preferred embodiment of the present invention, the selected transfer vehicle is capable of enhancing the stability of the mRNA contained therein. The liposome can allow the encapsulated mRNA to reach the target cell and/or may preferentially allow the encapsulated mRNA to reach the target cell, or alternatively limit the delivery of such mRNA to other sites or cells where the presence of the administered mRNA may be useless or undesirable. Furthermore, incorporating the mRNA into a transfer vehicle, such as for example, a cationic liposome, also facilitates the delivery of such mRNA into a target cell.

Ideally, liposomal transfer vehicles are prepared to encapsulate one or more desired mRNA such that the compositions demonstrate a high transfection efficiency and enhanced stability. While liposomes can facilitate introduction of nucleic acids into target cells, the addition of polycations (e.g., poly L-lysine and protamine), as a copolymer can facilitate, and in some instances markedly enhance the transfection efficiency of several types of cationic liposomes by 2-28 fold in a number of cell lines both in vitro and in vivo. (See N. J. Caplen, et al., *Gene Ther.* 1995; 2: 603; S. Li, et al., *Gene Ther.* 1997; 4, 891.)

Lipid Nanoparticles

In a preferred embodiment of the present invention, the transfer vehicle is formulated as a lipid nanoparticle. As used herein, the phrase “lipid nanoparticle” refers to a transfer vehicle comprising one or more lipids (e.g., cationic lipids, non-cationic lipids, and PEG-modified lipids). Preferably, the lipid nanoparticles are formulated to deliver one or more mRNA to one or more target cells. Examples of suitable lipids include, for example, the phosphatidyl compounds (e.g., phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides). Also contemplated is the use of polymers as transfer vehicles, whether alone or in combination with other transfer vehicles. Suitable polymers may include, for example, polyacrylates, polyalkylcyanoacrylates, polylactide, polylactide-polyglycolide copolymers, polycaprolactones, dextran, albumin, gelatin, alginate, collagen, chitosan, cyclodextrins, dendrimers and polyethylenimine. In one embodiment, the transfer vehicle is selected based upon its ability to facilitate the transfection of a mRNA to a target cell.

The invention contemplates the use of lipid nanoparticles as transfer vehicles comprising a cationic lipid to encapsulate and/or enhance the delivery of mRNA into the target cell that will act as a depot for protein production. As used herein, the phrase “cationic lipid” refers to any of a number of lipid species that carry a net positive charge at a selected pH, such as physiological pH. The contemplated lipid nanoparticles may be prepared by including multi-component lipid mixtures of varying ratios employing one or more cationic lipids, non-cationic lipids and PEG-modified lipids. Several cationic lipids have been described in the literature, many of which are commercially available.

Particularly suitable cationic lipids for use in the compositions and methods of the invention include those described in international patent publication WO 2010/053572, incorporated herein by reference, and most particularly, C12-200 described at paragraph [00225] of WO 2010/053572. In certain embodiments, the compositions and methods of the invention employ a lipid nanoparticles comprising an ionizable cationic lipid described in U.S. provisional patent application 61/617,468, filed Mar. 29, 2012 (incorporated herein by reference), such as, e.g., (15Z,18Z)-N,N-dimethyl-6-(9Z,12Z)-octadeca-9,12-dien-1-yl)tetracos-15,18-dien-1-amine (HGT5000), (15Z,18Z)-N,N-dimethyl-6-((9Z,12Z)-octadeca-9,12-dien-1-yl)tetracos-4,15,18-trien-1-amine (HGT5001), and (15Z,18Z)-N,N-dimethyl-6-((9Z,12Z)-octadeca-9,12-dien-1-yl)tetracos-5,15,18-trien-1-amine (HGT5002).

In some embodiments, the cationic lipid N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride or “DOTMA” is used. (Felgner et al. (*Proc. Nat'l Acad. Sci.* 84, 7413 (1987); U.S. Pat. No. 4,897,355). DOTMA can be formulated alone or can be combined with the neutral lipid, dioleoylphosphatidyl-ethanolamine or “DOPE” or other cationic or non-cationic lipids into a liposomal transfer vehicle or a lipid nanoparticle, and such liposomes can be used to enhance the delivery of nucleic acids into target cells. Other suitable cationic lipids include, for example, 5-carboxyspermylglycinedioctadecylamide or “DOGS,” 2,3-dioleoyloxy-N-[2(spermine-carboxamido)ethyl]-N,N-dimethyl-1-propanaminium or “DOSPA” (Behr et al. *Proc. Nat'l Acad. Sci.* 86, 6982 (1989); U.S. Pat. Nos. 5,171,678; 5,334,761), 1,2-Dioleoyl-3-Dimethylammonium-Propane or “DOD-AP”, 1,2-Dioleoyl-3-Trimethylammonium-Propane or “DOTAP”. Contemplated cationic lipids also include 1,2-distearoyloxy-N,N-dimethyl-3-aminopropane or “DSDMA”, 1,2-dioleoyloxy-N,N-dimethyl-3-aminopropane or “DODMA”, 1,2-dilinoleoyloxy-N,N-dimethyl-3-aminopropane or “DLinDMA”, 1,2-dilinoleoyloxy-N,N-dimethyl-3-aminopropane or “DLenDMA”, N-dioleoyl-N,N-dimethylammonium chloride or “DODAC”, N,N-distearoyl-N,N-dimethylammonium bromide or “DDAB”, N-(1,2-dimyristyloxypropyl-3-yl)-N,N-dimethyl-N-hydroxyethyl ammonium bromide or “DMRIE”, 3-dimethylamino-2-(cholest-5-en-3-beta-oxybutan-4-oxy)-1-(cis,cis-9,12-octadecadienoxy)propane or “CLinDMA”, 2-[5'-(cholest-5-en-3-beta-oxy)-3'-oxapentoxo]-3-dimethyl 1-1-(cis,cis-9',1-2'-octadecadienoxy)propane or “CpLinDMA”, N,N-dimethyl-3,4-dioleoyloxybenzylamine or “DMOBA”, 1,2-N,N'-dioleoylcarbonyl-3-dimethylaminopropane or “DOcarbDAP”, 2,3-Dilinoleoyloxy-N,N-dimethylpropylamine or “DLinDAP”, 1,2-N,N'-Dilinoleoylcarbonyl-3-dimethylaminopropane or “DLincarbDAP”, 1,2-Dilinoleoylcarbonyl-3-dimethylaminopropane or “DLinCDAP”, 2,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane or “DLin-K-DMA”, 2,2-dilinoleyl-4-dimethylaminoethyl[1,3]-dioxolane or “DLin-K-XTC2-DMA”, and 2-(2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)-1,3-dioxolan-4-yl)-N,N-dimethylethylamine (DLin-KC2-DMA)) (See, WO 2010/042877; Semple et al., *Nature Biotech.* 28:172-176 (2010)), or mixtures thereof. (Heyes, J., et al., *J Controlled Release* 107: 276-287 (2005); Morrissey, D V., et al., *Nat. Biotechnol.* 23(8): 1003-1007 (2005); PCT Publication WO2005/121348A1).


The use of cholesterol-based cationic lipids is also contemplated by the present invention. Such cholesterol-based cationic lipids can be used, either alone or in combination with other cationic or non-cationic lipids. Suitable cholesterol-based cationic lipids include, for example, DC-Chol

17


(N,N-dimethyl-N-ethylcarboxamidocholesterol), 1,4-bis(3-N-oleylamino-propyl)piperazine (Gao, et al. *Biochem. Biophys. Res. Comm.* 179, 280 (1991); Wolf et al. *BioTechniques* 23, 139 (1997); U.S. Pat. No. 5,744,335), or ICE.

In addition, several reagents are commercially available to enhance transfection efficacy. Suitable examples include LIPOFECTIN (DOTMA:DOPE) (Invitrogen, Carlsbad, Calif.), LIPOFECTAMINE (DOSPA:DOPE) (Invitrogen), LIPOFECTAMINE2000 (Invitrogen), FUGENE, TRANSFECTAM (DOGS), and EFFECTENE.

Also contemplated are cationic lipids such as the dialkylamino-based, imidazole-based, and guanidinium-based lipids. For example, certain embodiments are directed to a



composition comprising one or more imidazole-based cationic lipids, for example, the imidazole cholesterol ester or “ICE” lipid (3S,10R,13R,17R)-10,13-dimethyl-17-((R)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl 3-(1H-imidazol-4-yl)propanoate, as represented by structure (I) below. In a preferred embodiment, a transfer vehicle for delivery of mRNA may comprise one or more imidazole-based cationic lipids, for example, the imidazole cholesterol ester or “ICE” lipid (3S,10R,13R,17R)-10,13-dimethyl-17-((R)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl 3-(1H-imidazol-4-yl)propanoate, as represented by structure (I).



Without wishing to be bound by a particular theory, it is believed that the fusogenicity of the imidazole-based cationic lipid ICE is related to the endosomal disruption which is facilitated by the imidazole group, which has a lower pKa relative to traditional cationic lipids. The endosomal disruption in turn promotes osmotic swelling and the disruption of the liposomal membrane, followed by the transfection or intracellular release of the nucleic acid(s) contents loaded therein into the target cell.

The imidazole-based cationic lipids are also characterized by their reduced toxicity relative to other cationic lipids. The imidazole-based cationic lipids (e.g., ICE) may be used as the sole cationic lipid in the lipid nanoparticle, or alternatively may be combined with traditional cationic lipids, non-cationic lipids, and PEG-modified lipids. The cationic lipid may comprise a molar ratio of about 1% to about 90%, about 2% to about 70%, about 5% to about 50%, about 10%

18

to about 40% of the total lipid present in the transfer vehicle, or preferably about 20% to about 70% of the total lipid present in the transfer vehicle.

Similarly, certain embodiments are directed to lipid nanoparticles comprising the HGT4003 cationic lipid 2-((2,3-Bis((9Z,12Z)-octadeca-9,12-dien-1-yloxy)propyl)disulfanyl)-N,N-dimethylethanamine, as represented by structure (II) below, and as further described in U.S. Provisional Application No. 61/494,745, filed Jun. 8, 2011, the entire teachings of which are incorporated herein by reference in their entirety:

(I)

In other embodiments the compositions and methods described herein are directed to lipid nanoparticles comprising one or more cleavable lipids, such as, for example, one or more cationic lipids or compounds that comprise a cleavable disulfide (S—S) functional group (e.g., HGT4001, HGT4002, HGT4003, HGT4004 and HGT4005), as further described in U.S. Provisional Application No. 61/494,745, the entire teachings of which are incorporated herein by reference in their entirety.

The use of polyethylene glycol (PEG)-modified phospholipids and derivatized lipids such as derivatized ceramides (PEG-CER), including N-Octanoyl-Sphingosine-1-[Succinyl(Methoxy Polyethylene Glycol)-2000] (C8 PEG-2000 ceramide) is also contemplated by the present invention, either alone or preferably in combination with other lipids together which comprise the transfer vehicle (e.g., a lipid nanoparticle). Contemplated PEG-modified lipids include, but is not limited to, a polyethylene glycol chain of up to 5 kDa in length covalently attached to a lipid with alkyl chain(s) of C₅-C₂₀ length. The addition of such components may prevent complex aggregation and may also provide a means for increasing circulation lifetime and increasing the delivery of the lipid-nucleic acid composition to the target cell, (Klibanov et al. (1990) *FEBS Letters*, 268 (1): 235-237), or they may be selected to rapidly exchange out of the formulation in vivo (see U.S. Pat. No. 5,885,613). Particularly useful exchangeable lipids are PEG-ceramides having shorter acyl chains (e.g., C14 or C18). The PEG-modified phospholipid and derivatized lipids of the present invention may comprise a molar ratio from about 0% to about 20%, about 0.5% to about 20%, about 1% to about 15%, about 4% to about 10%, or about 2% of the total lipid present in the liposomal transfer vehicle.

The present invention also contemplates the use of non-cationic lipids. As used herein, the phrase “non-cationic lipid” refers to any neutral, zwitterionic or anionic lipid. As used herein, the phrase “anionic lipid” refers to any of a number of lipid species that carry a net negative charge at a selected pH, such as physiological pH. Non-cationic lipids include, but are not limited to, distearoylphosphatidylcholine (DSPC), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), dioleoylphosphatidylglycerol (DOPG), dipalmitoylphosphatidylglycerol (DPPG), dioleoylphosphatidylethanolamine (DOPE),

palmitoyloleoylphosphatidylcholine (POPC), palmitoyloleoyl-phosphatidylethanolamine (POPE), dioleoyl-phosphatidylethanolamine 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (DOPE-mal), dipalmitoyl phosphatidyl ethanolamine (DPPE), dimyristoylphosphoethanolamine (DMPE), distearoyl-phosphatidyl-ethanolamine (DSPE), 16-O-monomethyl PE, 16-O-dimethyl PE, 18-1-trans PE, 1-stearoyl-2-oleoyl-phosphatidylethanolamine (SOPE), cholesterol, or a mixture thereof. Such non-cationic lipids may be used alone, but are preferably used in combination with other excipients, for example, cationic lipids. When used in combination with a cationic lipid, the non-cationic lipid may comprise a molar ratio of 5% to about 90%, or preferably about 10% to about 70% of the total lipid present in the transfer vehicle.

Preferably, the transfer vehicle (e.g., a lipid nanoparticle) is prepared by combining multiple lipid and/or polymer components. For example, a transfer vehicle may be prepared using C12-200, DOPE, chol, DMG-PEG2K at a molar ratio of 40:30:25:5, or DODAP, DOPE, cholesterol, DMG-PEG2K at a molar ratio of 18:56:20:6, or HGT5000, DOPE, chol, DMG-PEG2K at a molar ratio of 40:20:35:5, or HGT5001, DOPE, chol, DMG-PEG2K at a molar ratio of 40:20:35:5. The selection of cationic lipids, non-cationic lipids and/or PEG-modified lipids which comprise the lipid nanoparticle, as well as the relative molar ratio of such lipids to each other, is based upon the characteristics of the selected lipid(s), the nature of the intended target cells, the characteristics of the mRNA to be delivered. Additional considerations include, for example, the saturation of the alkyl chain, as well as the size, charge, pH, pKa, fusogenicity and toxicity of the selected lipid(s). Thus the molar ratios may be adjusted accordingly. For example, in embodiments, the percentage of cationic lipid in the lipid nanoparticle may be greater than 10%, greater than 20%, greater than 30%, greater than 40%, greater than 50%, greater than 60%, or greater than 70%. The percentage of non-cationic lipid in the lipid nanoparticle may be greater than 5%, greater than 10%, greater than 20%, greater than 30%, or greater than 40%. The percentage of cholesterol in the lipid nanoparticle may be greater than 10%, greater than 20%, greater than 30%, or greater than 40%. The percentage of PEG-modified lipid in the lipid nanoparticle may be greater than 1%, greater than 2%, greater than 5%, greater than 10%, or greater than 20%.

In certain preferred embodiments, the lipid nanoparticles of the invention comprise at least one of the following cationic lipids: C12-200, DLin-KC2-DMA, DODAP, HGT4003, ICE, HGT5000, or HGT5001. In embodiments, the transfer vehicle comprises cholesterol and/or a PEG-modified lipid. In some embodiments, the transfer vehicles comprises DMG-PEG2K. In certain embodiments, the transfer vehicle comprises one of the following lipid formulations: C12-200, DOPE, chol, DMG-PEG2K; DODAP, DOPE, cholesterol, DMG-PEG2K; HGT5000, DOPE, chol, DMG-PEG2K, HGT5001, DOPE, chol, DMG-PEG2K.

The liposomal transfer vehicles for use in the compositions of the invention can be prepared by various techniques which are presently known in the art. Multi-lamellar vesicles (MLV) may be prepared conventional techniques, for example, by depositing a selected lipid on the inside wall of a suitable container or vessel by dissolving the lipid in an appropriate solvent, and then evaporating the solvent to leave a thin film on the inside of the vessel or by spray drying. An aqueous phase may then added to the vessel with a vortexing motion which results in the formation of MLVs. Uni-lamellar vesicles (ULV) can then be formed by homog-

enization, sonication or extrusion of the multi-lamellar vesicles. In addition, unilamellar vesicles can be formed by detergent removal techniques.

In certain embodiments of this invention, the compositions of the present invention comprise a transfer vehicle wherein the mRNA is associated on both the surface of the transfer vehicle and encapsulated within the same transfer vehicle. For example, during preparation of the compositions of the present invention, cationic liposomal transfer vehicles may associate with the mRNA through electrostatic interactions.

In certain embodiments, the compositions of the invention may be loaded with diagnostic radionuclide, fluorescent materials or other materials that are detectable in both in vitro and in vivo applications. For example, suitable diagnostic materials for use in the present invention may include Rhodamine-dioleoylphosphatidylethanolamine (Rh-PE), Green Fluorescent Protein mRNA (GFP mRNA), *Renilla* Luciferase mRNA and Firefly Luciferase mRNA.

Selection of the appropriate size of a liposomal transfer vehicle must take into consideration the site of the target cell or tissue and to some extent the application for which the liposome is being made. In some embodiments, it may be desirable to limit transfection of the mRNA to certain cells or tissues. For example, to target hepatocytes a liposomal transfer vehicle may be sized such that its dimensions are smaller than the fenestrations of the endothelial layer lining hepatic sinusoids in the liver; accordingly the liposomal transfer vehicle can readily penetrate such endothelial fenestrations to reach the target hepatocytes. Alternatively, a liposomal transfer vehicle may be sized such that the dimensions of the liposome are of a sufficient diameter to limit or expressly avoid distribution into certain cells or tissues. For example, a liposomal transfer vehicle may be sized such that its dimensions are larger than the fenestrations of the endothelial layer lining hepatic sinusoids to thereby limit distribution of the liposomal transfer vehicle to hepatocytes. Generally, the size of the transfer vehicle is within the range of about 25 to 250 nm, preferably less than about 250 nm, 175 nm, 150 nm, 125 nm, 100 nm, 75 nm, 50 nm, 25 nm or 10 nm.

A variety of alternative methods known in the art are available for sizing of a population of liposomal transfer vehicles. One such sizing method is described in U.S. Pat. No. 4,737,323, incorporated herein by reference. Sonicating a liposome suspension either by bath or probe sonication produces a progressive size reduction down to small ULV less than about 0.05 microns in diameter. Homogenization is another method that relies on shearing energy to fragment large liposomes into smaller ones. In a typical homogenization procedure, MLV are recirculated through a standard emulsion homogenizer until selected liposome sizes, typically between about 0.1 and 0.5 microns, are observed. The size of the liposomal vesicles may be determined by quasi-electric light scattering (QELS) as described in Bloomfield, *Ann. Rev. Biophys. Bioeng.*, 10:421-450 (1981), incorporated herein by reference. Average liposome diameter may be reduced by sonication of formed liposomes. Intermittent sonication cycles may be alternated with QELS assessment to guide efficient liposome synthesis.

Target Cells

As used herein, the term "target cell" refers to a cell or tissue to which a composition of the invention is to be directed or targeted. In some embodiments, the target cells are deficient in a protein or enzyme of interest. For example, where it is desired to deliver a nucleic acid to a hepatocyte, the hepatocyte represents the target cell. In some embodi-

ments, the compositions of the invention transfect the target cells on a discriminatory basis (i.e., do not transfect non-target cells). The compositions of the invention may also be prepared to preferentially target a variety of target cells, which include, but are not limited to, hepatocytes, epithelial cells, hematopoietic cells, epithelial cells, endothelial cells, lung cells, bone cells, stem cells, mesenchymal cells, neural cells (e.g., meninges, astrocytes, motor neurons, cells of the dorsal root ganglia and anterior horn motor neurons), photoreceptor cells (e.g., rods and cones), retinal pigmented epithelial cells, secretory cells, cardiac cells, adipocytes, vascular smooth muscle cells, cardiomyocytes, skeletal muscle cells, beta cells, pituitary cells, synovial lining cells, ovarian cells, testicular cells, fibroblasts, B cells, T cells, reticulocytes, leukocytes, granulocytes and tumor cells.

The compositions of the invention may be prepared to preferentially distribute to target cells such as in the heart, lungs, kidneys, liver, and spleen. In some embodiments, the compositions of the invention distribute into the cells of the liver to facilitate the delivery and the subsequent expression of the mRNA comprised therein by the cells of the liver (e.g., hepatocytes). The targeted hepatocytes may function as a biological "reservoir" or "depot" capable of producing, and systemically excreting a functional protein or enzyme. Accordingly, in one embodiment of the invention the liposomal transfer vehicle may target hepatocytes and/or preferentially distribute to the cells of the liver upon delivery. Following transfection of the target hepatocytes, the mRNA loaded in the liposomal vehicle are translated and a functional protein product is produced, excreted and systemically distributed. In other embodiments, cells other than hepatocytes (e.g., lung, spleen, heart, ocular, or cells of the central nervous system) can serve as a depot location for protein production.

In one embodiment, the compositions of the invention facilitate a subject's endogenous production of one or more functional proteins and/or enzymes, and in particular the production of proteins and/or enzymes which demonstrate less immunogenicity relative to their recombinantly-prepared counterparts. In a preferred embodiment of the present invention, the transfer vehicles comprise mRNA which encode a deficient protein or enzyme. Upon distribution of such compositions to the target tissues and the subsequent transfection of such target cells, the exogenous mRNA loaded into the liposomal transfer vehicle (e.g., a lipid nanoparticle) may be translated in vivo to produce a functional protein or enzyme encoded by the exogenously administered mRNA (e.g., a protein or enzyme in which the subject is deficient). Accordingly, the compositions of the present invention exploit a subject's ability to translate exogenously- or recombinantly-prepared mRNA to produce an endogenously-translated protein or enzyme, and thereby produce (and where applicable excrete) a functional protein or enzyme. The expressed or translated proteins or enzymes may also be characterized by the in vivo inclusion of native post-translational modifications which may often be absent in recombinantly-prepared proteins or enzymes, thereby further reducing the immunogenicity of the translated protein or enzyme.

The administration of mRNA encoding a deficient protein or enzyme avoids the need to deliver the nucleic acids to specific organelles within a target cell (e.g., mitochondria). Rather, upon transfection of a target cell and delivery of the nucleic acids to the cytoplasm of the target cell, the mRNA contents of a transfer vehicle may be translated and a functional protein or enzyme expressed.

The present invention also contemplates the discriminatory targeting of target cells and tissues by both passive and active targeting means. The phenomenon of passive targeting exploits the natural distributions patterns of a transfer vehicle in vivo without relying upon the use of additional excipients or means to enhance recognition of the transfer vehicle by target cells. For example, transfer vehicles which are subject to phagocytosis by the cells of the reticulo-endothelial system are likely to accumulate in the liver or spleen, and accordingly may provide means to passively direct the delivery of the compositions to such target cells.

Alternatively, the present invention contemplates active targeting, which involves the use of additional excipients, referred to herein as "targeting ligands" that may be bound (either covalently or non-covalently) to the transfer vehicle to encourage localization of such transfer vehicle at certain target cells or target tissues. For example, targeting may be mediated by the inclusion of one or more endogenous targeting ligands (e.g., apolipoprotein E) in or on the transfer vehicle to encourage distribution to the target cells or tissues. Recognition of the targeting ligand by the target tissues actively facilitates tissue distribution and cellular uptake of the transfer vehicle and/or its contents in the target cells and tissues (e.g., the inclusion of an apolipoprotein-E targeting ligand in or on the transfer vehicle encourages recognition and binding of the transfer vehicle to endogenous low density lipoprotein receptors expressed by hepatocytes). As provided herein, the composition can comprise a ligand capable of enhancing affinity of the composition to the target cell. Targeting ligands may be linked to the outer bilayer of the lipid particle during formulation or post-formulation. These methods are well known in the art. In addition, some lipid particle formulations may employ fusogenic polymers such as PEAA, hemagglutinin, other lipopeptides (see U.S. patent application Ser. Nos. 08/835,281, and 60/083,294, which are incorporated herein by reference) and other features useful for in vivo and/or intracellular delivery. In other some embodiments, the compositions of the present invention demonstrate improved transfection efficacies, and/or demonstrate enhanced selectivity towards target cells or tissues of interest. Contemplated therefore are compositions which comprise one or more ligands (e.g., peptides, aptamers, oligonucleotides, a vitamin or other molecules) that are capable of enhancing the affinity of the compositions and their nucleic acid contents for the target cells or tissues. Suitable ligands may optionally be bound or linked to the surface of the transfer vehicle. In some embodiments, the targeting ligand may span the surface of a transfer vehicle or be encapsulated within the transfer vehicle. Suitable ligands and are selected based upon their physical, chemical or biological properties (e.g., selective affinity and/or recognition of target cell surface markers or features.) Cell-specific target sites and their corresponding targeting ligand can vary widely. Suitable targeting ligands are selected such that the unique characteristics of a target cell are exploited, thus allowing the composition to discriminate between target and non-target cells. For example, compositions of the invention may include surface markers (e.g., apolipoprotein-B or apolipoprotein-E) that selectively enhance recognition of, or affinity to hepatocytes (e.g., by receptor-mediated recognition of and binding to such surface markers). Additionally, the use of galactose as a targeting ligand would be expected to direct the compositions of the present invention to parenchymal hepatocytes, or alternatively the use of mannose containing sugar residues as a targeting ligand would be expected to direct the compositions of the present invention to liver endothelial cells (e.g., mannose containing sugar

residues that may bind preferentially to the asialoglycoprotein receptor present in hepatocytes). (See Hillery A M, et al. "Drug Delivery and Targeting: For Pharmacists and Pharmaceutical Scientists" (2002) Taylor & Francis, Inc.) The presentation of such targeting ligands that have been conjugated to moieties present in the transfer vehicle (e.g., a lipid nanoparticle) therefore facilitate recognition and uptake of the compositions of the present invention in target cells and tissues. Examples of suitable targeting ligands include one or more peptides, proteins, aptamers, vitamins and oligonucleotides.

Application and Administration

As used herein, the term "subject" refers to any animal (e.g., a mammal), including, but not limited to, humans, non-human primates, rodents, and the like, to which the compositions and methods of the present invention are administered. Typically, the terms "subject" and "patient" are used interchangeably herein in reference to a human subject.

The compositions and methods of the invention provide for the delivery of mRNA to treat a number of disorders. In particular, the compositions and methods of the present invention are suitable for the treatment of diseases or disorders relating to the deficiency of proteins and/or enzymes that are excreted or secreted by the target cell into the surrounding extracellular fluid (e.g., mRNA encoding hormones and neurotransmitters). In embodiments the disease may involve a defect or deficiency in a secreted protein (e.g. Fabry disease, or ALS). In certain embodiments, the disease may not be caused by a defect or deficit in a secreted protein, but may benefit from providing a secreted protein. For example, the symptoms of a disease may be improved by providing the compositions of the invention (e.g. cystic fibrosis). Disorders for which the present invention are useful include, but are not limited to, disorders such as Huntington's Disease; Parkinson's Disease; muscular dystrophies (such as, e.g. Duchenne and Becker); hemophilia diseases (such as, e.g., hemophilia B (FIX), hemophilia A (FVIII); SMN1-related spinal muscular atrophy (SMA); amyotrophic lateral sclerosis (ALS); GALT-related galactosemia; Cystic Fibrosis (CF); SLC3A1-related disorders including cystinuria; COL4A5-related disorders including Alport syndrome; galactocerebrosidase deficiencies; X-linked adrenoleukodystrophy and adrenomyeloneuropathy; Friedreich's ataxia; Pelizaeus-Merzbacher disease; TSC1 and TSC2-related tuberous sclerosis; Sanfilippo B syndrome (MPS IIIB); CTNS-related cystinosis; the FMR1-related disorders which include Fragile X syndrome, Fragile X-Associated Tremor/Ataxia Syndrome and Fragile X Premature Ovarian Failure Syndrome; Prader-Willi syndrome; hereditary hemorrhagic telangiectasia (AT); Niemann-Pick disease Type C1; the neuronal ceroid lipofuscinosis-related diseases including Juvenile Neuronal Ceroid Lipofuscinosis (JNCL), Juvenile Batten disease, Santavuori-Haltia disease, Jansky-Bielschowsky disease, and PTT-1 and TPP1 deficiencies; EIF2B1, EIF2B2, EIF2B3, EIF2B4 and EIF2B5-related childhood ataxia with central nervous system hypomyelination/vanishing white matter; CACNA1A and CACNB4-related Episodic Ataxia Type 2; the MECP2-related disorders including Classic Rett Syndrome, MECP2-related Severe Neonatal Encephalopathy and PPM-X Syndrome; CDKL5-related Atypical Rett Syndrome; Kennedy's disease (SBMA); Notch-3 related cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL); SCN1A and SCN1B-related seizure disorders; the Polymerase G-related disorders which include Alpers-Huttenlocher syndrome, POLG-related sen-

sory ataxic neuropathy, dysarthria, and ophthalmoparesis, and autosomal dominant and recessive progressive external ophthalmoplegia with mitochondrial DNA deletions; X-Linked adrenal hypoplasia; X-linked agammaglobulinemia; Wilson's disease; and Fabry Disease. In one embodiment, the nucleic acids, and in particular mRNA, of the invention may encode functional proteins or enzymes that are secreted into extracellular space. For example, the secreted proteins include clotting factors, components of the complement pathway, cytokines, chemokines, chemoattractants, protein hormones (e.g. EGF, PDF), protein components of serum, antibodies, secretable toll-like receptors, and others. In some embodiments, the compositions of the present invention may include mRNA encoding erythropoietin, α 1-antitrypsin, carboxypeptidase N or human growth hormone.

In embodiments, the invention encodes a secreted protein that is made up of subunits that are encoded by more than one gene. For example, the secreted protein may be a heterodimer, wherein each chain or subunit of the is encoded by a separate gene. It is possible that more than one mRNA molecule is delivered in the transfer vehicle and the mRNA encodes separate subunit of the secreted protein. Alternatively, a single mRNA may be engineered to encode more than one subunit (e.g. in the case of a single-chain Fv antibody). In certain embodiments, separate mRNA molecules encoding the individual subunits may be administered in separate transfer vehicles. In one embodiment, the mRNA may encode full length antibodies (both heavy and light chains of the variable and constant regions) or fragments of antibodies (e.g. Fab, Fv, or a single chain Fv (scFv) to confer immunity to a subject. While one embodiment of the present invention relates to methods and compositions useful for conferring immunity to a subject (e.g., via the translation of mRNA encoding functional antibodies), the inventions disclosed herein and contemplated hereby are broadly applicable. In an alternative embodiment the compositions of the present invention encode antibodies that may be used to transiently or chronically effect a functional response in subjects. For example, the mRNA of the present invention may encode a functional monoclonal or polyclonal antibody, which upon translation and secretion from target cell may be useful for targeting and/or inactivating a biological target (e.g., a stimulatory cytokine such as tumor necrosis factor). Similarly, the mRNA nucleic acids of the present invention may encode, for example, functional anti-nephritic factor antibodies useful for the treatment of membranoproliferative glomerulonephritis type II or acute hemolytic uremic syndrome, or alternatively may encode anti-vascular endothelial growth factor (VEGF) antibodies useful for the treatment of VEGF-mediated diseases, such as cancer. In other embodiments, the secreted protein is a cytokine or other secreted protein comprised of more than one subunit (e.g. IL-12, or IL-23).

The compositions of the invention can be administered to a subject. In some embodiments, the composition is formulated in combination with one or more additional nucleic acids, carriers, targeting ligands or stabilizing reagents, or in pharmacological compositions where it is mixed with suitable excipients. For example, in one embodiment, the compositions of the invention may be prepared to deliver mRNA encoding two or more distinct proteins or enzymes. Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition.

A wide range of molecules that can exert pharmaceutical or therapeutic effects can be delivered into target cells using

compositions and methods of the invention. The molecules can be organic or inorganic. Organic molecules can be peptides, proteins, carbohydrates, lipids, sterols, nucleic acids (including peptide nucleic acids), or any combination thereof. A formulation for delivery into target cells can comprise more than one type of molecule, for example, two different nucleotide sequences, or a protein, an enzyme or a steroid.

The compositions of the present invention may be administered and dosed in accordance with current medical practice, taking into account the clinical condition of the subject, the site and method of administration, the scheduling of administration, the subject's age, sex, body weight and other factors relevant to clinicians of ordinary skill in the art. The "effective amount" for the purposes herein may be determined by such relevant considerations as are known to those of ordinary skill in experimental clinical research, pharmacological, clinical and medical arts. In some embodiments, the amount administered is effective to achieve at least some stabilization, improvement or elimination of symptoms and other indicators as are selected as appropriate measures of disease progress, regression or improvement by those of skill in the art. For example, a suitable amount and dosing regimen is one that causes at least transient protein production.

Suitable routes of administration include, for example, oral, rectal, vaginal, transmucosal, pulmonary including intratracheal or inhaled, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, the compositions of the invention may be administered in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into a targeted tissue, preferably in a sustained release formulation. Local delivery can be affected in various ways, depending on the tissue to be targeted. For example, aerosols containing compositions of the present invention can be inhaled (for nasal, tracheal, or bronchial delivery); compositions of the present invention can be injected into the site of injury, disease manifestation, or pain, for example; compositions can be provided in lozenges for oral, tracheal, or esophageal application; can be supplied in liquid, tablet or capsule form for administration to the stomach or intestines, can be supplied in suppository form for rectal or vaginal application; or can even be delivered to the eye by use of creams, drops, or even injection. Formulations containing compositions of the present invention complexed with therapeutic molecules or ligands can even be surgically administered, for example in association with a polymer or other structure or substance that can allow the compositions to diffuse from the site of implantation to surrounding cells. Alternatively, they can be applied surgically without the use of polymers or supports.

In one embodiment, the compositions of the invention are formulated such that they are suitable for extended-release of the mRNA contained therein. Such extended-release compositions may be conveniently administered to a subject at extended dosing intervals. For example, in one embodiment, the compositions of the present invention are administered to a subject twice day, daily or every other day. In a preferred embodiment, the compositions of the present invention are administered to a subject twice a week, once a week, every ten days, every two weeks, every three weeks, or more preferably every four weeks, once a month, every six weeks, every eight weeks, every other month, every three

months, every four months, every six months, every eight months, every nine months or annually. Also contemplated are compositions and liposomal vehicles which are formulated for depot administration (e.g., intramuscularly, subcutaneously, intravitreally) to either deliver or release a mRNA over extended periods of time. Preferably, the extended-release means employed are combined with modifications made to the mRNA to enhance stability.

Also contemplated herein are lyophilized pharmaceutical compositions comprising one or more of the liposomal nanoparticles disclosed herein and related methods for the use of such lyophilized compositions as disclosed for example, in U.S. Provisional Application No. 61/494,882, filed Jun. 8, 2011, the teachings of which are incorporated herein by reference in their entirety. For example, lyophilized pharmaceutical compositions according to the invention may be reconstituted prior to administration or can be reconstituted in vivo. For example, a lyophilized pharmaceutical composition can be formulated in an appropriate dosage form (e.g., an intradermal dosage form such as a disk, rod or membrane) and administered such that the dosage form is rehydrated over time in vivo by the individual's bodily fluids.

While certain compounds, compositions and methods of the present invention have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds of the invention and are not intended to limit the same. Each of the publications, reference materials, accession numbers and the like referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference in their entirety.

The articles "a" and "an" as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to include the plural referents. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention also includes embodiments in which more than one, or the entire group members are present in, employed in, or otherwise relevant to a given product or process. Furthermore, it is to be understood that the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, descriptive terms, etc., from one or more of the listed claims is introduced into another claim dependent on the same base claim (or, as relevant, any other claim) unless otherwise indicated or unless it would be evident to one of ordinary skill in the art that a contradiction or inconsistency would arise. Where elements are presented as lists, (e.g., in Markush group or similar format) it is to be understood that each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements, features, etc., certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements, features, etc. For purposes of simplicity those embodiments have not in every case been specifically set forth in so many words herein. It should also be understood that any embodiment or aspect of the invention can be

explicitly excluded from the claims, regardless of whether the specific exclusion is recited in the specification. The publications and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

EXAMPLES

Example 1: Protein Production Depot Via Intravenous Delivery of Polynucleotide Compositions

Messenger RNA

Human erythropoietin (EPO) (SEQ ID NO: 3; FIG. 3), human alpha-galactosidase (GLA) (SEQ ID NO: 4; FIG. 4), human alpha-1 antitrypsin (A1AT) (SEQ ID NO: 5; FIG. 5), and human factor IX (FIX) (SEQ ID NO: 6; FIG. 6) were synthesized by in vitro transcription from a plasmid DNA template encoding the gene, which was followed by the addition of a 5' cap structure (Cap1) (Fechter & Brownlee, *J. Gen. Virology* 86:1239-1249 (2005)) and a 3' poly(A) tail of approximately 200 nucleotides in length as determined by gel electrophoresis. 5' and 3' untranslated regions were present in each mRNA product in the following examples and are defined by SEQ ID NOs: 1 and 2 (FIG. 1 and FIG. 2) respectively.

Lipid Nanoparticle Formulations

Formulation 1: Aliquots of 50 mg/mL ethanolic solutions of C12-200, DOPE, Chol and DMG-PEG2K (40:30:25:5) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C.

Formulation 2: Aliquots of 50 mg/mL ethanolic solutions of DODAP, DOPE, cholesterol and DMG-PEG2K (18:56:20:6) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of EPO mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C. Final concentration=1.35 mg/mL EPO mRNA (encapsulated). Z_{ave} =75.9 nm ($Dv_{(50)}$ =57.3 nm; $Dv_{(90)}$ =92.1 nm).

Formulation 3: Aliquots of 50 mg/mL ethanolic solutions of HGT4003, DOPE, cholesterol and DMG-PEG2K (50:25:20:5) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C.

Formulation 4: Aliquots of 50 mg/mL ethanolic solutions of ICE, DOPE and DMG-PEG2K (70:25:5) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20%

ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C.

Formulation 5: Aliquots of 50 mg/mL ethanolic solutions of HGT5000, DOPE, cholesterol and DMG-PEG2K (40:20:35:5) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of EPO mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C. Final concentration=1.82 mg/mL EPO mRNA (encapsulated). Z_{ave} =105.6 nm ($Dv_{(50)}$ =53.7 nm; $Dv_{(90)}$ =157 nm).

Formulation 6: Aliquots of 50 mg/mL ethanolic solutions of HGT5001, DOPE, cholesterol and DMG-PEG2K (40:20:35:5) were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of EPO mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1×PBS (pH 7.4), concentrated and stored at 2-8° C.

Analysis of Protein Produced Via Intravenously Delivered mRNA-Loaded Nanoparticles Injection Protocol

Studies were performed using male CD-1 mice of approximately 6-8 weeks of age at the beginning of each experiment, unless otherwise indicated. Samples were introduced by a single bolus tail-vein injection of an equivalent total dose of 30-200 micrograms of encapsulated mRNA. Mice were sacrificed and perfused with saline at the designated time points.

Isolation of Organ Tissues for Analysis

The liver and spleen of each mouse was harvested, apportioned into three parts, and stored in either 10% neutral buffered formalin or snap-frozen and stored at -80° C. for analysis.

Isolation of Serum for Analysis

All animals were euthanized by CO₂ asphyxiation 48 hours post dose administration ($\pm 5\%$) followed by thoracotomy and terminal cardiac blood collection. Whole blood (maximal obtainable volume) was collected via cardiac puncture on euthanized animals into serum separator tubes, allowed to clot at room temperature for at least 30 minutes, centrifuged at 22° C. $\pm 5^\circ$ C. at 9300 g for 10 minutes, and the serum extracted. For interim blood collections, approximately 40-50 μ L of whole blood was collected via facial vein puncture or tail snip. Samples collected from non treatment animals were used as a baseline for comparison to study animals.

Enzyme-Linked Immunosorbent Assay (ELISA) Analysis
EPO ELISA: Quantification of EPO protein was performed following procedures reported for human EPO ELISA kit (Quantikine IVD, R&D Systems, Catalog #Dep-00). Positive controls employed consisted of ultrapure and tissue culture grade recombinant human erythropoietin protein (R&D Systems, Catalog #286-EP and 287-TC, respectively). Detection was monitored via absorption (450 nm) on a Molecular Device Flex Station instrument.

GLA ELISA: Standard ELISA procedures were followed employing sheep anti-Alpha-galactosidase G-188 IgG as the capture antibody with rabbit anti-Alpha-galactosidase TK-88 IgG as the secondary (detection) antibody (Shire Human Genetic Therapies). Horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG was used for activation of

the 3,3',5,5'-tetramethylbenzidine (TMB) substrate solution. The reaction was quenched using 2N H₂SO₄ after 20 minutes. Detection was monitored via absorption (450 nm) on a Molecular Device Flex Station instrument. Untreated mouse serum and human Alpha-galactosidase protein were used as negative and positive controls, respectively.

FIX ELISA: Quantification of FIX protein was performed following procedures reported for human FIX ELISA kit (AssayMax, Assay Pro, Catalog #EF1009-1).

A1AT ELISA: Quantification of A1AT protein was performed following procedures reported for human A1AT ELISA kit (Innovative Research, Catalog #IRAPKT015).

Western Blot Analysis

(EPO): Western blot analyses were performed using an anti-hEPO antibody (R&D Systems #MAB2871) and ultra-pure human EPO protein (R&D Systems #286-EP) as the control.

Results

The work described in this example demonstrates the use of mRNA-encapsulated lipid nanoparticles as a depot source for the production of protein. Such a depot effect can be achieved in multiple sites within the body (i.e., liver, kidney, spleen, and muscle). Measurement of the desired exogenous-based protein derived from messenger RNA delivered via liposomal nanoparticles was achieved and quantified, and the secretion of protein from a depot using human erythropoietin (hEPO), human alpha-galactosidase (hGLA), human alpha-1 antitrypsin (hA1AT), and human Factor IX (hFIX) mRNA was demonstrated.

1A. In Vivo Human EPO Protein Production Results

The production of hEPO protein was demonstrated with various lipid nanoparticle formulations. Of four different cationic lipid systems, C12-200-based lipid nanoparticles produced the highest quantity of hEPO protein after four hours post intravenous administration as measured by ELISA (FIG. 7). This formulation (Formulation 1) resulted in 18.3 ug/mL hEPO protein secreted into the bloodstream. Normal hEPO protein levels in serum for human are 3.3-16.6 mIU/mL (NCCLS Document C28-P; Vol. 12, No. 2). Based on a specific activity of 120,000 IU/mg of EPO protein, that yields a quantity of 27.5-138 pg/mL hEPO protein in normal human individuals. Therefore, a single 30 ug dose of a C12-200-based cationic lipid formulation encapsulating hEPO mRNA yielded an increase in respective protein of over 100,000-fold physiological levels.

Of the lipid systems tested, the DODAP-based lipid nanoparticle formulation was the least effective. However, the observed quantity of human EPO protein derived from delivery via a DODAP-based lipid nanoparticle encapsulating EPO mRNA was 4.1 ng/mL, which is still greater than 30-fold over normal physiological levels of EPO protein (Table 1).

TABLE 1

Raw values of secreted hEPO protein for various cationic lipid-based nanoparticle systems as measured via ELISA analysis (as depicted in FIG. 8). Doses are based on encapsulated hEPO mRNA. Values of protein are depicted as nanogram of human EPO protein per milliliter of serum. Hematocrit changes are based on comparison of pre-bleed (Day -1) and Day 10.			
Cationic/Ionizable Lipid Component	Dose of Encapsulated mRNA (ug)	Secreted Human EPO Protein (ng/mL)	Increase in Hematocrit (%)
C12-200	30	18,306	15.0
HGT4003	150	164	0.0

TABLE 1-continued

Raw values of secreted hEPO protein for various cationic lipid-based nanoparticle systems as measured via ELISA analysis (as depicted in FIG. 8). Doses are based on encapsulated hEPO mRNA. Values of protein are depicted as nanogram of human EPO protein per milliliter of serum. Hematocrit changes are based on comparison of pre-bleed (Day -1) and Day 10.			
Cationic/Ionizable Lipid Component	Dose of Encapsulated mRNA (ug)	Secreted Human EPO Protein (ng/mL)	Increase in Hematocrit (%)
ICE	100	56.2	0.0
DODAP	200	4.1	0.0

In addition, the resulting protein was tested to determine if it was active and functioned properly. In the case of mRNA replacement therapy (MRT) employing hEPO mRNA, hematocrit changes were monitored over a ten day period for five different lipid nanoparticle formulations (FIG. 8, Table 1) to evaluate protein activity. During this time period, two of the five formulations demonstrated an increase in hematocrit ($\geq 15\%$), which is indicative of active hEPO protein being produced from such systems.

In another experiment, hematocrit changes were monitored over a 15-day period (FIG. 9, Table 2). The lipid nanoparticle formulation (Formulation 1) was administered either as a single 30 μ g dose, or as three smaller 10 μ g doses injected on day 1, day 3 and day 5. Similarly, Formulation 2 was administered as 3 doses of 50 μ g on day 1, day 3, and day 5. C12-200 produced a significant increase in hematocrit. Overall an increase of up to ~25% change was observed, which is indicative of active human EPO protein being produced from such systems.

TABLE 2

Hematocrit levels of each group over a 15 day observation period (FIG. 9). Mice were either dosed as a single injection, or three injections, every other day. N = 4 mice per group.					
Test Article	Dose (μ g/animal)	Hct Levels Mean (%) \pm SEM			
		Day -4	Day 7	Day 10	Day 15 ^a
C12-200	30 (single dose)	50.8 \pm 1.8	58.3 \pm 3.3	62.8 \pm 1.3	59.9 \pm 3.3
C12-200	30 (over 3 doses)	52.2 \pm 0.5	55.3 \pm 2.3	63.3 \pm 1.6	62.3 \pm 1.9
DODAP	150 (over 3 doses)	54.8 \pm 1.7	53.5 \pm 1.6	54.2 \pm 3.3	54.0 \pm 0.3

Hct = hematocrit;

SEM = standard error of the mean.

^aBlood samples were collected into non-heparinized hematocrit tubes.

1B. In Vivo Human GLA Protein Production Results

A second exogenous-based protein system was explored to demonstrate the "depot effect" when employing mRNA-loaded lipid nanoparticles. Animals were injected intravenously with a single 30 microgram dose of encapsulated human alpha-galactosidase (hGLA) mRNA using a C12-200-based lipid nanoparticle system and sacrificed after six hours (Formulation 1). Quantification of secreted hGLA protein was performed via ELISA. Untreated mouse serum and human Alpha-galactosidase protein were used as controls. Detection of human alpha-galactosidase protein was monitored over a 48 hour period.

Measurable levels of hGLA protein were observed throughout the time course of the experiment with a maxi-

imum level of 2.0 ug/mL hGLA protein at six hours (FIG. 10). Table 3 lists the specific quantities of hGLA found in the serum. Normal activity in healthy human males has been reported to be approximately 3.05 nanomol/hr/mL. The

TABLE 3

Raw values of secreted hGLA protein over time as measured via ELISA analysis (as depicted in FIG. 10). Values are depicted as nanogram of hGLA protein per milliliter of serum. N = 4 mice per group.	
Time Post-Administration (hr)	Secreted Human GLA Protein (ng/mL)
6	2,038
12	1,815
24	414
48	86.2

In addition, the half-life of Alpha-galactosidase when administered at 0.2 mg/kg is approximately 108 minutes. Production of GLA protein via the “depot effect” when administering GLA mRNA-loaded lipid nanoparticles shows a substantial increase in blood residence time when compared to direct injection of the naked recombinant protein. As described above, significant quantities of protein are present after 48 hours.

The activity profile of the α -galactosidase protein produced from GLA mRNA-loaded lipid nanoparticles was measured as a function of 4-methylumbelliferyl- α -D-galactopyranoside (4-MU- α -gal) metabolism. As shown in FIG. 11, the protein produced from these nanoparticle systems is quite active and reflective of the levels of protein available

(FIG. 12, Table 3). AUC comparisons of mRNA therapy-based hGLA production versus enzyme replacement therapy (ERT) in mice and humans show a 182-fold and 30-fold increase, respectively (Table 4).

TABLE 4

Comparison of C_{max} and AUC_{inf} values in Fabry patients post-IV dosing 0.2 mg/kg of Alpha-galactosidase (pharmacological dose) with those in mice post-IV dosing Alpha-galactosidase and GLA mRNA.

Test Article	Description	Dose (mg/kg)	C_{max} (U/mL)	AUC_{inf} (hr.U/mL)	n	
Fabry ^a Patient	α -GAL Protein	Transplant	0.2	3478	11	
		Dialysis	0.2	3887	6	
		Non-ESRD ^b	0.2	3710	18	
Mouse	α -GAL Protein (MM1)	Athymic nude	0.04	3807	3	
	α -GAL Protein (MM2)	Athymic nude	0.04	3705	3	
Mouse	α -GAL mRNA	mouse	0.95	5885 ($C_{at 6hr}$) ^c	109428	6

^aData were from a published paper (Gregory M. Pastores et al. Safety and Pharmacokinetics of hGLA in patients with Fabry disease and end-stage renal disease. *Nephrol Dial Transplant* (2007) 22: 1920-1925.

^bnon-end-stage renal disease.

^c α -Galactosidase activity at 6 hours after dosing (the earliest time point tested in the study).

activity for Alpha-galactosidase, a recombinant human alpha-galactosidase protein, 3.56×10^6 nanomol/hr/mg. Analysis of these values yields a quantity of approximately 856 pg/mL of hGLA protein in normal healthy male individuals. The quantity of 2.0 ug/mL hGLA protein observed after six hours when dosing a hGLA mRNA-loaded lipid nanoparticle is over 2300-fold greater than normal physiological levels. Further, after 48 hours, one can still detect appreciable levels of hGLA protein (86.2 ng/mL). This level is representative of almost 100-fold greater quantities of hGLA protein over physiological amounts still present at 48 hours.

The ability of mRNA encapsulated lipid nanoparticles to target organs which can act as a depot for the production of a desired protein has been demonstrated. The levels of secreted protein observed have been several orders of magnitude above normal physiological levels. This “depot effect” is repeatable. FIG. 12 shows again that robust protein production is observed upon dosing wild type (CD-1) mice with a single 30 ug dose of hGLA mRNA-loaded in C12-200-based lipid nanoparticles (Formulation 1). In this experiment, hGLA levels were evaluated over a 72 hour period. A maximum average of 4.0 ug human hGLA protein/mL serum is detected six hours post-administration. Based on a value of ~1 ng/mL hGLA protein for normal physiological levels, hGLA MRT provides roughly 4000-fold higher protein levels. As before, hGLA protein could be detected out to 48 hr post-administration (FIG. 12).

An analysis of tissues isolated from this same experiment provided insight into the distribution of hGLA protein in hGLA MRT-treated mice (FIG. 13). Supraphysiological levels of hGLA protein were detected in the liver, spleen and kidneys of all mice treated with a maximum observed between 12 and 24 hour post-administration. Detectable levels of MRT-derived protein could be observed three days after a single injection of hGLA-loaded lipid nanoparticles.

In addition, the production of hGLA upon administration of hGLA mRNA loaded C12-200 nanoparticles was shown to exhibit a dose response in the serum (FIG. 14A), as well as in the liver (FIG. 14B).

One inherent characteristic of lipid nanoparticle-mediated mRNA replacement therapy would be the pharmacokinetic profile of the respective protein produced. For example, ERT-based treatment of mice employing Alpha-galactosidase results in a plasma half-life of approximately 100 minutes. In contrast, MRT-derived alpha-galactosidase has a blood residence time of approximately 72 hrs with a peak time of 6 hours. This allows for much greater exposure for organs to participate in possible continuous uptake of the desired protein. A comparison of PK profiles is shown in FIG. 15 and demonstrates the stark difference in clearance rates and ultimately a major shift in area under the curve (AUC) can be achieved via MRT-based treatment.

In a separate experiment, hGLA MRT was applied to a mouse disease model, hGLA KO mice (Fabry mice). A 0.33 mg/kg dose of hGLA mRNA-loaded C12-200-based lipid nanoparticles (Formulation 1) was administered to female KO mice as a single, intravenous injection. Substantial quantities of MRT-derived hGLA protein were produced with a peak at 6 hr (~560 ng/mL serum) which is approximately 600-fold higher than normal physiological levels. Further, hGLA protein was still detectable 72 hr post-administration (FIG. 16).

Quantification of MRT-derived GLA protein in vital organs demonstrated substantial accumulation as shown in FIG. 17. A comparison of observed MRT-derived hGLA protein to reported normal physiological levels that are found in key organs is plotted (normal levels plotted as dashed lines). While levels of protein at 24 hours are higher than at 72 hours post-administration, the levels of hGLA protein detected in the liver, kidney, spleen and hearts of the treated Fabry mice are equivalent to wild type levels. For example, 3.1 ng hGLA protein/mg tissue were found in the kidneys of treated mice 3 days after a single MRT treatment.

In a subsequent experiment, a comparison of ERT-based Alpha-galactosidase treatment versus hGLA MRT-based treatment of male Fabry KO mice was conducted. A single, intravenous dose of 1.0 mg/kg was given for each therapy and the mice were sacrificed one week post-administration. Serum levels of hGLA protein were monitored at 6 hr and 1 week post-injection. Liver, kidney, spleen, and heart were analyzed for hGLA protein accumulation one week post-administration. In addition to the biodistribution analyses, a measure of efficacy was determined via measurement of globotriaosylceramide (Gb3) and lyso-Gb3 reductions in the kidney and heart. FIG. 18 shows the serum levels of hGLA protein after treatment of either Alpha-galactosidase or GLA mRNA loaded lipid nanoparticles (Formulation 1) in male Fabry mice. Serum samples were analyzed at 6 hr and 1 week post-administration. A robust signal was detected for MRT-treated mice after 6 hours, with hGLA protein serum levels of ~4.0 ug/mL. In contrast, there was no detectable Alpha-galactosidase remaining in the bloodstream at this time.

The Fabry mice in this experiment were sacrificed one week after the initial injection and the organs were harvested and analyzed (liver, kidney, spleen, heart). FIG. 19 shows a comparison of human GLA protein found in each respective organ after either hGLA MRT or Alpha-galactosidase ERT treatment. Levels correspond to hGLA present one week post-administration. hGLA protein was detected in all organs analyzed. For example, MRT-treated mice resulted in hGLA protein accumulation in the kidney of 2.42 ng hGLA protein/mg protein, while Alpha-galactosidase-treated mice had only residual levels (0.37 ng/mg protein). This corresponds to a ~6.5-fold higher level of hGLA protein when treated via hGLA MRT. Upon analysis of the heart, 11.5 ng hGLA protein/mg protein was found for the MRT-treated cohort as compared to only 1.0 ng/mg protein Alpha-galactosidase. This corresponds to an ~11-fold higher accumulation in the heart for hGLA MRT-treated mice over ERT-based therapies.

In addition to the biodistribution analyses conducted, evaluations of efficacy were determined via measurement of globotriaosylceramide (Gb3) and lyso-Gb3 levels in key organs. A direct comparison of Gb3 reduction after a single, intravenous 1.0 mg/kg GLA MRT treatment as compared to a Alpha-galactosidase ERT-based therapy of an equivalent dose yielded a sizeable difference in levels of Gb3 in the kidneys as well as heart. For example, Gb3 levels for GLA

MRT versus Alpha-galactosidase yielded reductions of 60.2% vs 26.8%, respectively (FIG. 20). Further, Gb3 levels in the heart were reduced by 92.1% vs 66.9% for MRT and Alpha-galactosidase, respectively (FIG. 21).

A second relevant biomarker for measurement of efficacy is lyso-Gb3. GLA MRT reduced lyso-Gb3 more efficiently than Alpha-galactosidase as well in the kidneys and heart (FIG. 20 and FIG. 21, respectively). In particular, MRT-treated Fabry mice demonstrated reductions of lyso-Gb3 of 86.1% and 87.9% in the kidneys and heart as compared to Alpha-galactosidase-treated mice yielding a decrease of 47.8% and 61.3%, respectively.

The results with for hGLA in C12-200 based lipid nanoparticles extend to other lipid nanoparticle formulations. For example, hGLA mRNA loaded into HGT4003 (Formulation 3) or HGT5000-based (Formulation 5) lipid nanoparticles administered as a single dose IV result in production of hGLA at 24 hours post administration (FIG. 22). The production of hGLA exhibited a dose response. Similarly, hGLA production was observed at 6 hours and 24 hours after administration of hGLA mRNA loaded into HGT5001-based (Formulation 6) lipid nanoparticles administered as a single dose IV. hGLA production was observed in the serum (FIG. 23A), as well as in organs (FIG. 23B).

Overall, mRNA replacement therapy applied as a depot for protein production produces large quantities of active, functionally therapeutic protein at supraphysiological levels. This method has been demonstrated to yield a sustained circulation half-life of the desired protein and this MRT-derived protein is highly efficacious for therapy as demonstrated with alpha-galactosidase enzyme in Fabry mice.

1C. In Vivo Human FIX Protein Production Results

Studies were performed administering Factor IX (FIX) mRNA-loaded lipid nanoparticles in wild type mice (CD-1) and determining FIX protein that is secreted into the bloodstream. Upon intravenous injection of a single dose of 30 ug C12-200-based (C12-200:DOPE:Chol:PEG at a ratio of 40:30:25:5) FIX mRNA-loaded lipid nanoparticles (dose based on encapsulated mRNA) (Formulation 1), a robust protein production was observed (FIG. 24).

A pharmacokinetic analysis over 72 hours showed MRT-derived FIX protein could be detected at all timepoints tested (FIG. 24). The peak serum concentration was observed at 24 hr post-injection with a value of ~3 ug (2995±738 ng/mL) FIX protein/mL serum. This represents another successful example of the depot effect.

1D. In Vivo Human A1AT Protein Production Results

Studies were performed administering alpha-1-antitrypsin (A1AT) mRNA-loaded lipid nanoparticles in wild type mice (CD-1) and determining A1AT protein that is secreted into the bloodstream. Upon intravenous injection of a single dose of 30 ug C12-200-based A1AT mRNA-loaded lipid nanoparticles (dose based on encapsulated mRNA) (Formulation 1), a robust protein production was observed (FIG. 25).

As depicted in FIG. 25, detectable levels of human A1AT protein derived from A1AT MRT could be observed over a 24 hour time period post-administration. A maximum serum level of ~48 ug A1AT protein/mL serum was detected 12 hours after injection.

Example 2: Protein Production Depot Via Pulmonary Delivery of Polynucleotide Compositions

65 Injection Protocol

All studies were performed using female CD-1 or BALB/C mice of approximately 7-10 weeks of age at the

beginning of each experiment. Test articles were introduced via a single intratracheal aerosolized administration. Mice were sacrificed and perfused with saline at the designated time points. The lungs of each mouse were harvested, apportioned into two parts, and stored in either 10% neutral buffered formalin or snap-frozen and stored at -80°C . for analysis. Serum was isolated as described in Example 1. EPO ELISA: as described in Example 1.

Results

The depot effect can be achieved via pulmonary delivery (e.g. intranasal, intratracheal, nebulization). Measurement of the desired exogenous-based protein derived from messenger RNA delivered via nanoparticle systems was achieved and quantified.

The production of human EPO protein via hEPO mRNA-loaded lipid nanoparticles was tested in CD-1 mice via a single intratracheal administration (MicroSprayer®). Several formulations were tested using various cationic lipids (Formulations 1, 5, 6). All formulations resulted in high encapsulation of human EPO mRNA. Upon administration, animals were sacrificed six hours post-administration and the lungs as well as serum were harvested.

Human EPO protein was detected at the site of administration (lungs) upon treatment via aerosol delivery. Analysis of the serum six hours post-administration showed detectable amounts of hEPO protein in circulation. These data (shown in FIG. 26) demonstrate the ability of the lung to act as a “depot” for the production (and secretion) of hEPO protein.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 140

<212> TYPE: RNA

<213> ORGANISM: Human cytomegalovirus

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(3)

<223> OTHER INFORMATION: This region may or may not be present

<400> SEQUENCE: 1

```
ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac      60
cgauccagcc uccgcggccg ggaacggugc auuggaacgc ggauuccccg ugccaagagu     120
gacucaccgu ccuugacacg                                     140
```

<210> SEQ ID NO 2

<211> LENGTH: 100

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```
cggguggcau cccugugacc ccucccagu gccucuccug gcccuggaag uugccacucc      60
agugcccacc agccuugucc uaauaaaauu aaguugcauc                                     100
```

<210> SEQ ID NO 3

<211> LENGTH: 582

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```
augggggugc acgaaugucc ugccuggcug ugguucucc ugucccugcu gucgcucccu      60
cugggccucc caguccuggg cgccccacca cgccucaucu gugacagccg aguccuggag     120
agguaccucu uggaggccaa ggaggccgag aauaucacga cgggcugugc ugaacacugc     180
agcuugaaug agaauaucac ugucccagac accaaaguua auuucuaugc cuggaagagg     240
auggaggucg ggcagcaggc cguagaaguc uggcagggcc uggcccugcu gucgggaagcu     300
guccugcggg gccaggcccu guuggucaac ucuucccagc cgugggagcc ccugcagcug     360
cauguggaua aagccgucag ugccuucgc agccucacca cucugcuucg ggcucuggga     420
gccagaagg aagccaucuc cccuccagau gcggccucag cugcuccacu ccgaacaauc     480
acugcugaca cuuuccgcaa acucuuccga gucuacucca auuuccuccg gggaaagcug     540
aagcuguaca caggggaggc cugcaggaca ggggacagau ga                               582
```

-continued

```

<210> SEQ ID NO 4
<211> LENGTH: 1290
<212> TYPE: RNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4
augcagcuga ggaaccaga acuacaucug ggcugcgcgc uugcgcuucg cuuccuggcc 60
cucguuuccu gggacauccc uggggcuaga gcacuggaca auggauuggc aaggacgccu 120
accaugggcu ggcugcacug ggagcgcuuc augugcaacc uugacugcca ggaagagcca 180
gauuccugca ucagugagaa gcucuucaug gagauggcag agcucauggu cucagaaggc 240
uggaaggau gagguauga guaccucugc auugaugacu guuggauggc uccccaaaga 300
gauucagaag gcagacuca ggcagaccu cagcgcuuuc cucaugggau ucgccagcua 360
gcuaauuaug uucacagcaa aggacugaag cuagggauuu augcagaugu uggaaauaaa 420
accugcgcag gcuucccugg gaguuuugga uacuacgaca uugaugccca gaccuuugcu 480
gacuggggag uagaucugcu aaaauuugau gguuguuacu gugacaguuu ggaaaauuug 540
gcagaugguu auaagcacau guccuuggcc cugaauagga cuggcagaag cauuguguac 600
uccugugagu ggccucuua uauguggccc uuucaaaagc ccauuuauac agaaauccga 660
caguacugca aucacuggcg aaauuuugcu gacauugaug auuccuggaa aaguauaaag 720
aguaucuugg acuggacauc uuuuaaccag gagagaauug uugauguugc uggaccaggg 780
gguuggaaug acccagauau guuagugauu ggcaacuug gccucagcug gaaucagcaa 840
guaacucaga uggcccucug ggcuaucaug gcugcuccu uauucauguc uaaugaccuc 900
cgacacauca gcccucaagc caaagcucuc cuucaggaua aggacguauu ugccaaucau 960
caggaccccu ugggcaagca aggguaaccag cuuagacagg gagacaacu ugaagugugg 1020
gaacgaccuc ucucaggcuu agccugggcu guagcuauga uaaaccggca ggagauuggu 1080
ggaccucgcu cuuauaccu cgcaguugcu ucccugggua aaggaguggc cuguauuccu 1140
gccugcuuca ucacacagcu ccucccugug aaaaggaagc uaggguuuca ugaauaggacu 1200
ucaagguuaa gaagucacau aaaucccaca ggcacuguuu ugcucagcu agaaaauaca 1260
augcagaugu cauuuuuaga cuuacuuaa 1290

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1257
<212> TYPE: RNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5
augccgucuu cugucucgug gggcauccuc cugcuggcag gccugugcug ccugguccu 60
gucucccugg cugaggaucc ccagggagau gcugcccaga agacagauac aucccaccu 120
gaucaggauc acccaaccu caacaagauc accccaacc uggcugaguu cgccuucagc 180
cuauaccgcc agcuggcaca ccaguccaac agcaccaua ucuucucuc cccagugagc 240
aucguacag ccuuugcaau gcucucccug gggaccaagg cugacacuca cgaugaauc 300
cuggagggcc ugaauuucaa ccucacggag auuccggagg cucagaucua ugaaggcuuc 360
caggaacucc uccguaccu caaccagcca gacagccagc uccagcugac caccggcau 420
ggccuguucc ucagcgagg ccugaagcua guggauaagu uuuuggagga uguuuuuuag 480
uuguaccacu cagaagccu cacugucaac uucggggaca ccgaagaggc caagaaacag 540
aucaacgauu acguggagaa gguacucaa gggaaaauug uggauuuggu caaggagcu 600

```

-continued

```

gacagagaca caguuuuugc ucuggugaau uacauucuuc uuaaaggcaa augggagaga 660
cccuuugaag ucaaggacac cgaggaagag gacuuccacg uggaccaggu gaccaccgug 720
aaggugccua ugaugaagcg uuuaggcaug uuuuacauc agcacuguaa gaagcugucc 780
agcugggugc ugcugaugaa auaccugggc aaugccaccg ccaucuucu ccugccugau 840
gaggggaaac uacagcaccu ggaaaaugaa cucaccacg auaucaucac caaguuccug 900
gaaaaugaag acagaagguc ugccagcuua cauuuaccca aacuguccau uacuggaacc 960
uaugaucuga agagcguccu gggucaacug ggcaucacua aggucuucag caauggggcu 1020
gaccucuccg gggucacaga ggaggcacc cugaagcucu ccaaggccgu gcauaaggcu 1080
gugcugacca ucgacgagaa agggacugaa gcugcugggg ccauguuuuu agaggccaua 1140
cccaugucua uccccccga ggucaaguuc acaaacccu uugucuucu aaugauugaa 1200
caaaaauacca agucuccccu cuucauggga aaagugguga auccaccca aaaaaua 1257

```

<210> SEQ ID NO 6

<211> LENGTH: 1386

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

augcagcgcg ugaacaugau cauggcagaa ucaccaggcc ucaucaccau cugccuuuuu 60
ggauaucuac ucagugcuga auguacaguu uuucuugauc augaaaacgc caacaaaauu 120
cugaggcgga gaaggaggua uaaucaggu aaauuggaag aguuuguuca agggaaccuu 180
gagagagaau guauggaaga aaaguguagu uuugaagaag cacgagaagu uuuugaaaac 240
acugaaagaa caacugaauu uuggaagcag uauguugaug gagaucagug ugaguccaau 300
ccauguuuua auggcggcag uugcaaggau gacauuaau ccuaugaug uugguguccc 360
uuuggauuug aaggaaagaa cugugaauua gauguaacau guaacaauaa gaauggcaga 420
ugcgagcagu uuuguaaaaa uagugcugau aacaaggugg uuugcuccug uacugaggga 480
uacgacuug cagaaaacca gaaguccugu gaaccagcag ugccauuucc auguggaaga 540
guuucuguuu cacaaacuuc uaagcucacc cgugcugagg cuguuuuucc ugauguggac 600
uauguaaaau cuacugaagc ugaaaccauu uuggauaaca ucacuaaag cacccaauca 660
uuuaaugacu ucacucgggu uguuggugga gaagaugcca aaccagguca auucccuugg 720
cagguuguuu ugaauuguaa aguugaugca uucuguggag gcucuaucgu uaaugaaaaa 780
uggauuguaa cugcugccca cuguguugaa acugguguaa aaauuacagu ugucgcaggu 840
gaacauaaua uugaggagac agaacaucac gagcaaaagc gaaaugugau ucgaauuuuu 900
ccucaccaca acuacaaugc agcuauuuau aaguacaacc augacaauugc ccuucuggaa 960
cuggacgaac ccuagugcu aaacagcuac guuacaccua uuugcauugc ugacaaggaa 1020
uacacgaaca ucuuccuca aauuggaucu ggcuauguaa guggcugggg aagagucuuc 1080
caciaaaggga gaucagcuuu aguucuucag uaccuuagag uuccacuugu ugaccgagcc 1140
acaugucuuc gaucuaaaa guucaccauc uauaacaaca uguucugugc uggcuuccau 1200
gaaggaggua gagauucaug ucaaggagau agugggggac cccauguuac ugaaguggaa 1260
gggaccaguu ucuuaacugg aauuuuagc uggggugaag agugugcaau gaaaggcaaa 1320
uauggaaauu auaccaaggu aucccggau gucaacugga uuaaggaaaa acaaaagcuc 1380
acuuuaa 1386

```

-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 90
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 7

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      60
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

```

<210> SEQ ID NO 8
<211> LENGTH: 200
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 8

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      60
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     120
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     180
aaaaaaaaa aaaaaaaaaa

```

```

<210> SEQ ID NO 9
<211> LENGTH: 300
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      60
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     120
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     180
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     240
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     300

```

```

<210> SEQ ID NO 10
<211> LENGTH: 400
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      60
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     120
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     180
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     240
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     300
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa     360
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

```

<210> SEQ ID NO 11
<211> LENGTH: 500

```

-continued

```

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 11

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      60
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      120
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      180
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      240
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      300
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      360
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      420
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      480
aaaaaaaaa aaaaaaaaaa                                     500
    
```

The invention claimed is:

1. A method for delivery of messenger RNA (mRNA) for in vivo production of a protein or a peptide comprising administering to a subject a composition comprising an mRNA that encodes the protein or the peptide, wherein the mRNA is encapsulated within a lipid nanoparticle and wherein the administering of the composition results in the expression of the protein or the peptide encoded by the mRNA, wherein the lipid nanoparticle comprises (i) a PEG-modified lipid at a molar ratio of greater than 1% of the total lipids in the lipid nanoparticle, (ii) a cationic lipid at a molar ratio of greater than 10% of the total lipids in the lipid nanoparticle, (iii) a non-cationic lipid at a molar ratio of greater than 5% of total lipids in the lipid nanoparticle, and (iv) cholesterol at greater than 10% of the total lipids in the lipid nanoparticle wherein the non-cationic lipid comprises DOPE or DSPC.

2. The method of claim 1, wherein the non-cationic lipid comprises DSPC.

3. The method of claim 2, wherein the lipid nanoparticle comprises PEG-modified lipids at a molar ratio of 1% to 20% of the total lipids.

4. The method of claim 3, wherein the lipid nanoparticle comprises non-cationic lipids at a molar ratio of 5% to 90% of the total lipids.

5. The method of claim 1, wherein the non-cationic lipid comprises DOPE.

6. The method of claim 5, wherein one or more PEG-modified lipids comprise DMG-PEG-2000.

7. The method of claim 1, wherein the composition is administered systemically.

8. The method of claim 1, wherein the composition is administered intramuscularly.

9. The method of claim 1, wherein the composition is administered subcutaneously.

10. The method of claim 1, wherein the administering of the composition results in the expression of the protein or the peptide encoded by the mRNA that is detectable in a target tissue at least 48 hours after administration.

11. The method of claim 1, wherein the mRNA is modified.

12. The method of claim 11, wherein the mRNA comprises pseudouridines.

13. The method of claim 1, wherein the mRNA comprises a 3' untranslated region.

14. The method of claim 1, wherein the mRNA comprises a cap structure.

15. The method of claim 1, wherein the mRNA comprises a poly A tail.

16. The method of claim 1, wherein the lipid nanoparticle has a size of less than 150 nm.

17. The method of claim 16, wherein the lipid nanoparticle has a size of less than 125 nm.

18. The method of claim 17, wherein the lipid nanoparticle has a size of less than 100 nm.

19. The method of claim 1, wherein the administering of the composition confers immunity to a subject.

20. The method of claim 1, wherein the target tissue comprises muscle cells.

21. The method of claim 20, wherein the muscles cells are skeletal muscle cells, or vascular smooth muscle cells.

* * * * *